arXiv:1208.6166v1 [math.AP] 30 Aug 2012

Construction of transmutation operators and hyperbolic pseudoanalytic functions

Vladislav V. Kravchenko and Sergii M. Torba

Abstract

A representation for integral kernels of Delsarte transmutation operators is obtained in the form of a functional series with the exact formulae for the terms of the series. It is based on the application of hyperbolic pseudoanalytic function theory and recent results on mapping properties of the transmutation operators.

The kernel K_1 of the transmutation operator relating $A=-\frac{d^2}{dx^2}+q_1(x)$ and $B=-\frac{d^2}{dx^2}$ results to be one of the complex components of a bicomplex-valued hyperbolic pseudoanalytic function satisfying a Vekua-type hyperbolic equation of a special form. The other component of the pseudoanalytic function is the kernel of the transmutation operator relating $C=-\frac{d^2}{dx^2}+q_2(x)$ and B where q_2 is obtained from q_1 by a Darboux transformation. We prove the expansion theorem and a Runge-type theorem for this special hyperbolic Vekua equation and using several known results from hyperbolic pseudoanalytic function theory together with the recently discovered mapping properties of the transmutation operators obtain the new representation for their kernels. Several examples are given. Moreover, based on the presented results an approach for numerical computation of the transmutation kernels is proposed.

1. *Introduction*

The notion of a transmutation operator relating two linear differential operators was introduced in 1938 by J. Delsarte [**12**] and nowadays represents a widely used tool in the theory of linear differential equations (see, e.g., [**1**], [**7**], [**32**], [**34**], [**42**], [**44**]). Very often in literature the transmutation operators are called the transformation operators. Here we keep the original term introduced by Delsarte and Lions [**13**]. It is well known that under certain regularity conditions the transmutation operator transmuting the operator $A=-\frac{d^2}{dx^2}+q(x)$ into $B=-\frac{d^2}{dx^2}$ is a Volterra integral operator with good properties. Its integral kernel can be obtained as a solution of a certain Goursat problem for the Klein-Gordon equation with a variable coefficient. There exist very few examples of the transmutation kernels available in a closed form (see [**26**]).

In the present work we obtain a representation for the kernels of the transmutation operators for regular Sturm-Liouville operators (with complex valued coefficients) in the form of a functional series with the exact formulae for the terms of the series. The result is based on several new observations. We use our recent result on the construction of the kernel of the transmutation operator corresponding to a Darboux associated Schrödinger operator [**26**] to find out that a bicomplex-valued function whose one complex component is the transmutation kernel $K_1(x,t)$ (for a Schrödinger operator $\frac{d^2}{dx^2}-q_1(x)$, $q_1\in C[-b,b]$) and the other complex component is $K_2(x,t)$ (for a Schrödinger operator $\frac{d^2}{dx^2}-q_2(x)$, with q_2 obtained from q_1 by a

---

2000 *Mathematics Subject Classification* 30B60, 30G20, 34A25, 35A35, 35C10 (primary), 30B10, 34L16, 34A45, 35L10, 41A50, 47N20, 65N99 (secondary).

Research was supported by CONACYT, Mexico. Research of second named author was supported by DFFD, Ukraine (GP/F32/030) and by SNSF, Switzerland (JRP IZ73Z0 of SCOPES 2009–2012).

Darboux transformation) is a solution of a certain hyperbolic Vekua equation of a special form (for the theory of elliptic Vekua equations see [**45**] as well as [**2**] and [**23**]).

In spite of a recent progress reported in [**20**], [**23**], [**25**], [**29**] the theory of hyperbolic Vekua equations is considerably less developed than the theory of classical (elliptic) Vekua equations. For example, as was shown in [**25**] (see also [**23**]) the construction of so-called formal powers (solutions of the Vekua equation generalizing the usual complex powers $(z-z_0)^n$) can be performed using the definitions completely analogous to those introduced by L. Bers [**2**]. Nevertheless no expansion theorem nor a result on the completeness of the obtained system (a Runge-type theorem) of hyperbolic formal powers was available. In this work we apply the results from [**25**] for constructing the formal powers for the hyperbolic Vekua equation arising in relation with the kernels K_1 and K_2 as well as the observation from [**6**] establishing that they are the result of the transmutation of the usual powers of the hyperbolic variable. We obtain an expansion theorem and prove a completeness result. Moreover, we obtain explicit formulae for the expansion coefficients which leads to the functional series representation for the kernel K_1 (and also for K_2). We give several examples of application of this result including a numerical computation. Finally, we propose an alternative method for approximate construction of the transmutation kernel based on the same hyperbolic formal powers but instead of the expansion theorem using the obtained completeness result.

The paper is structured as follows. In Section 2 we introduce the hyperbolic and bicomplex numbers, the Vekua equation one of the solutions of which is the bicomplex function $K_1-\mathbf{j}K_2$. We present several properties of that Vekua equation, construct an infinite system of its solutions called formal powers, explain its relation to generalized wave polynomials and to spectral parameter power series (SPPS) representation for solutions of Sturm-Liouville equations. We consider a Goursat problem for the hyperbolic Cauchy-Riemann system and obtain certain related results important for what follows. In Section 3 we recall some long known facts about the transmutation operators and give some recent and new results concerning the relation between K_1 and K_2, the commutation and the mapping properties of the corresponding transmutation operators. In Section 4 we introduce an operator transforming the boundary data of the Goursat problem for the wave equation into the boundary data of the Goursat problem for the operator $\Box-q(x)$. Using its properties we obtain a result about a functional series representation for solutions of the equation $(\Box-q)\,u=0$ in terms of generalized wave polynomials. In Section 5 we introduce a transmutation operator relating the hyperbolic Cauchy-Riemann operator with the interesting for us hyperbolic Vekua operator. The main observation here is that this transmutation operator possessing important boundedness properties maps usual powers of the hyperbolic variable z into the formal powers as well as the usual derivatives with respect to z into the generalized derivatives in the sense of L. Bers. Here we mainly use the ideas from [**5**] where similar observations were made in the elliptic setting. Section 6 is dedicated to the expansion and the Runge-type theorems corresponding to the special Vekua equation under consideration. We show that a generalized Taylor formula for its solutions is valid under certain conditions on their Goursat data. The formula involves the formal powers constructed and studied in preceding sections. The obtained Runge-type theorem establishes that even when the conditions required for the validity of the Taylor expansion are not fulfilled any solution of the studied hyperbolic Vekua equation can be approximated arbitrarily closely by a formal polynomial. In Section 7 we show that K_1 and K_2 are necessarily the conjugate components of a solution of the Vekua equation. Using this we give explicit formulae expressing K_2 in terms of K_1 and vice versa. Section 8 contains the main result of this work - the representation of the transmutation kernel in terms of a generalized Taylor series in formal powers. For the corresponding expansion coefficients we obtain both a recurrent and a direct exact formulae. Several examples are given illustrating their application. Finally, in Section 9 we show how the obtained results can be used for numerical computation of the transmutation kernels. Here together with the approach based on the generalized Taylor expansion theorem we propose a method based on the Runge-type theorem. When sufficiently many derivatives of the potential taken in the origin are not available this second method is more convenient. A developed numerical example illustrates its excellent computational performance.

2. *Elements of hyperbolic pseudoanalytic function theory*

2.1. *Hyperbolic and bicomplex numbers*

By $\mathbf{j}$ we denote the hyperbolic imaginary unit: $\mathbf{j}^2=1$ and consider the algebra of hyperbolic numbers, also called duplex numbers (see, e.g., [**43**])

$$\mathbb{D}:=\left\{c=a+b\mathbf{j}\ :\ \mathbf{j}^2=1,\ a,b\in\mathbb{R}\right\}\cong \mathrm{Cl}_{\mathbb{R}}(0,1).$$

The algebra $\mathbb{D}$ is commutative and contains zero divisors. Additional information on the hyperbolic numbers can be found in [**31**], [**36**] and [**43**].

We will consider the variable $z=x+t\mathbf{j}$ where x and t are real variables and the corresponding differential operators

$$\partial_z=\frac{1}{2}\left(\partial_x+\mathbf{j}\partial_t\right)\text{ and }\partial_{\bar z}=\frac{1}{2}\left(\partial_x-\mathbf{j}\partial_t\right).$$

As in the case of complex numbers, we have $\partial_{\bar z}z=0$ which explains the choice of the minus sign in the definition of $\partial_{\bar z}$.

Let $\mathbb{B}$ denote the algebra of bicomplex numbers which can be defined as follows

$$\mathbb{B}:=\left\{w=u+v\mathbf{j}\ :\ u,v\in\mathbb{C}\right\}$$

and the complex imaginary unit $\mathbf{i}$ commutes with $\mathbf{j}$. More on bicomplex numbers can be found, e.g., in [**4**], [**8**], [**23**] and [**41**].

The conjugate of a bicomplex number w with respect to $\mathbf{j}$ we denote by $\overline{w}$, i.e., $\overline{w}=u-v\mathbf{j}$. The corresponding conjugation operator is denoted by C, $Cw=\overline{w}$.

$\mathbb{C}$-valued functions will be also called scalar. For the scalar components of w we introduce the notations

$$\mathcal{R}(w)=u=\frac{1}{2}(w+\overline{w})\quad\text{and}\quad\mathcal{I}(w)=v=\frac{1}{2\mathbf{j}}(w-\overline{w}).$$

In order not to overload the text by excessively many different notations we will use the notation $\mathcal{R}$ and $\mathcal{I}$ also in the operational sense as projection operators projecting $w\in\mathbb{B}$ onto the respective scalar components, $\mathcal{R}=\frac{1}{2}(I+C)$ and $\mathcal{I}=\frac{1}{2\mathbf{j}}(I-C)$ where I is the identity operator.

It is convenient to introduce the pair of idempotents $P^+=\frac{1}{2}(1+\mathbf{j})$ and $P^-=\frac{1}{2}(1-\mathbf{j})$ satisfying $\left(P^\pm\right)^2=P^\pm$ and $P^+P^-=P^-P^+=0$. Then for any $w\in\mathbb{B}$ there exist the unique numbers w^+, $w^-\in\mathbb{C}$ such that $w=P^+w^++P^-w^-$ which are related with the components of w in the following way

$$w^\pm=\mathcal{R}\left(w\right)\pm\mathcal{I}(w). \tag{2.1}$$

A nonzero element $w\in\mathbb{B}$ belongs to the set of zero divisors $\sigma(\mathbb{B})$ iff $w=P^+w^+$ or $w=P^-w^-$.

We will use the following norm (see [**4**]) in $\mathbb{B}$,

$$|w|=\frac{1}{2}\left(\left|w^+\right|_{\mathbb{C}}+\left|w^-\right|_{\mathbb{C}}\right), \tag{2.2}$$

where $\left|\cdot\right|_{\mathbb{C}}$ is the usual norm in $\mathbb{C}$.

2.2. *Hyperbolic Vekua equation*

We will consider functions $w : \mathbb{D} \to \mathbb{B}$ and Vekua-type equations of the form

$$\partial_{\bar{z}} w = aw + b\overline{w} \tag{2.3}$$

where a and b are bicomplex functions of the variable $z \in \mathbb{D}$. In [**20**], [**23**], [**25**] it was shown that many results from pseudoanalytic function theory [**2**] remain valid in the hyperbolic situation. We will not give here definitions and properties corresponding to the general Vekua equation (2.3) referring the reader to [**23**]. Instead, as we are interested in a very special form of the coefficients a and b we restrict ourselves to that particular case and obtain several results, such as the expansion theorem and a Runge-type theorem which are still unavailable in a general situation. Thus, let us consider the following hyperbolic Vekua equation

$$\partial_{\bar{z}} W - \frac{\partial_{\bar{z}} f}{f} \overline{W} = 0 \tag{2.4}$$

where f is a scalar, non-vanishing function in the domain of interest. Vekua equations of this form, i.e., with $a \equiv 0$ and b being a logarithmic derivative of a scalar function are called Vekua equations of the main type or main Vekua equations. They arise from the factorization of the stationary Schrödinger operator (in the elliptic case) [**21**], [**23**] and of the variable mass Klein-Gordon operator (in the hyperbolic case) [**25**], [**23**]. Moreover, for the purposes of the present paper it is sufficient to consider the case when f depends only on x and hence (2.4) can be written also in the form

$$\partial_{\bar{z}} W - \frac{f'(x)}{2f(x)} \overline{W} = 0. \tag{2.5}$$

It is equivalent to the system

$$f \partial_x \left(\frac{1}{f} u \right) = \partial_t v, \tag{2.6}$$

$$\frac{1}{f} \partial_x (fv) = \partial_t u, \tag{2.7}$$

where u, v: $\mathbb{R}^2 \to \mathbb{C}$ and $W = u + v\mathbf{j}$. Introducing the notation $\varphi = u/f$ and $\psi = fv$ one can rewrite this system also in the following form

$$\partial_t \varphi = \frac{1}{f^2} \partial_x \psi \quad \text{and} \quad \partial_x \varphi = \frac{1}{f^2} \partial_t \psi. \tag{2.8}$$

In the case when all the involved functions were real valued this system was studied in [**37**], [**38**] and later on in [**25**], [**23**].

Equation (2.4) admits a corresponding generating pair $F = f$ and $G = \mathbf{j}/f$. We recall that a generating pair corresponding to a Vekua equation is a pair of its solutions F and G, independent in the sense that an arbitrary function ω, in this case $\mathbb{B}$-valued, can be represented in the form

$$\omega = \varphi F + \psi G \tag{2.9}$$

where φ and ψ are scalar functions. That is, F and G should satisfy the condition $\mathcal{I}(\overline{F}G) \neq 0$ everywhere in the domain of interest. In relation with the generating pair $(F, G) = (f, \mathbf{j}/f)$ observe that the scalar functions φ and ψ satisfy (2.8) iff the $\mathbb{B}$-valued function $\omega = \varphi f + \psi \mathbf{j}/f$ is a solution of (2.4).

The knowledge of a generating pair allows one to define the (F, G)-derivative in the sense of L. Bers. For a function ω written in the form (2.9) its (F, G)-derivative has the form

$$\overset{\circ}{\omega} := \frac{d_{(F,G)}\omega}{dz} := (\partial_z \varphi) F + (\partial_z \psi) G.$$

Whenever W satisfies (2.4) its (F,G)-derivative (with $(F,G)=(f,\mathbf{j}/f)$) is a solution of the Vekua equation [**23**]

$$\partial_{\bar{z}} w + \frac{\partial_z f}{f}\overline{w} = 0. \tag{2.10}$$

Notice that for $f=f(x)$ this equation can be written as follows

$$\partial_{\bar{z}} w + \frac{f'(x)}{2f(x)}\overline{w} = 0$$

or

$$\partial_{\bar{z}} w - \frac{\left(f^{-1}(x)\right)'}{2f^{-1}(x)}\overline{w} = 0. \tag{2.11}$$

Thus, the coefficient in the equation has the same structure as in (2.5) but the role of f is played by $1/f$. Hence a generating pair for (2.11) can be chosen as $(F_1,G_1)=(1/f,f\mathbf{j})$. The second derivative of a solution of (2.5) can be defined in the form $\overset{\circ\circ}{W} = \overset{\circ}{w}$ where $w = \overset{\circ}{W} = \varphi_1 F_1 + \psi_1 G_1 = \varphi_1/f + \psi_1 f\mathbf{j}$, φ_1 and ψ_1 are scalar functions, and $\overset{\circ}{w} = (\partial_z\varphi_1)/f + (\partial_z\psi_1) f\mathbf{j}$. Obviously, the next generating pair can be chosen again as $(F_2,G_2)=(f,\mathbf{j}/f)$ and in this way the derivative of any order in the sense of Bers can be defined. The derivative of n-th order we denote as $W^{[n]}$. In the theory of pseudoanalytic functions such a sequence of generating pairs is known as a periodic generating sequence (of the period two). We have $(F_n,G_n)=(f,\mathbf{j}/f)$ when n is even, and $(F_n,G_n)=(1/f,f\mathbf{j})$ when n is odd.

For what follows it is important to observe that if W is a solution of (2.5), its n-th derivative in the sense of Bers (if exists) has an especially simple form

$$W^{[n]} = \mathbf{j}^n \partial_t^n W. \tag{2.12}$$

Indeed, for $W=\varphi f + \psi\mathbf{j}/f$ we have $\overset{\circ}{W} = (\partial_z\varphi) f + (\partial_z\psi)\mathbf{j}/f$ where φ and ψ are solutions of (2.8). Consider

$$\partial_z\varphi = \frac{1}{2}\left(\partial_x\varphi + \mathbf{j}\partial_t\varphi\right) = \frac{1}{2}\left(\frac{1}{f^2}\partial_t\psi + \mathbf{j}\partial_t\varphi\right) = \frac{\mathbf{j}}{2f}\partial_t\left(\varphi f + \psi\frac{\mathbf{j}}{f}\right) = \frac{\mathbf{j}}{2f}\partial_t W$$

and similarly,

$$\partial_z\psi = \frac{1}{2}\left(\partial_x\psi + \mathbf{j}\partial_t\psi\right) = \frac{1}{2}\left(f^2\partial_t\varphi + \mathbf{j}\partial_t\psi\right) = \frac{f}{2}\partial_t\left(\varphi f + \psi\frac{\mathbf{j}}{f}\right) = \frac{f}{2}\partial_t W.$$

Thus, $\overset{\circ}{W} = \mathbf{j}\partial_t W$. As this reasoning does not change if f is substituted everywhere by $1/f$ we obtain that $\overset{\circ\circ}{W} = \mathbf{j}\partial_t\overset{\circ}{W} = \mathbf{j}^2\partial_t^2 W$ and hence (2.12) is valid both for odd and for even values of n.

Below we extend some of the results on the relationship between hyperbolic pseudoanalytic functions and solutions of the Klein-Gordon equations to the bicomplex case. Since all the proofs are essentially the same, we refer the reader to [**23**, Chapter 13], [**25**] for details. The operator $\partial_{\bar{z}}$ applied to a scalar function φ can be regarded as a kind of gradient. If $\partial_{\bar{z}}\varphi = \Phi$, where Φ is a $\mathbb{B}$-valued function defined on a simply connected domain Ω with $\Phi_1 = \mathcal{R}(\Phi)$ and $\Phi_2 = \mathcal{I}(\Phi)$ such that

$$\partial_t\Phi_1 + \partial_x\Phi_2 = 0, \quad \forall\,(x,t)\in\Omega,$$

then we can construct φ up to an arbitrary complex constant c. Indeed, let $\Gamma\subset\Omega$ be a rectifiable curve leading from (x_0,t_0) to (x,t), then the integral

$$\overline{A}_h[\Phi](x,t) := 2\left(\int_\Gamma \Phi_1\,dx - \Phi_2\,dt\right) \tag{2.13}$$

is path-independent, and all $\mathbb{C}$-valued solutions φ of the equation $\partial_{\bar z}\varphi = \Phi$ in Ω have the form $\varphi(x,t) = \overline{A}_h[\Phi(x,t)] + c$ where c is an arbitrary complex constant. In other words the operator $\overline{A}_h$ denotes the well known operation for reconstructing the potential function from its gradient. In Section 7 we need a particular case of (2.13) when Γ is the path consisting from 2 segments, first going from (x_0,t_0) to (x,t_0) and second going to (x,t). Assuming that this path belongs to the domain of interest Ω, in such case formula (2.13) reads as follows

$$\overline{A}_h[\Phi](x,t) = 2\left(\int_{x_0}^{x} \Phi_1(\eta,t_0)\,d\eta - \int_{t_0}^{t} \Phi_2(x,\xi)\,d\xi\right). \tag{2.14}$$

The following theorems explain relationship between the main Vekua equation and variable mass Klein-Gordon equations. We use the notation $\square := \partial_x^2 - \partial_t^2$.

Theorem 2.1 [**25**]. *Let $f = f(x)$ be a non-vanishing scalar function and W be a solution of* (2.5)*. Then $u = \mathcal{R}(W)$ is a solution of the equation*

$$\big(\square - q_1(x)\big)u = 0, \quad \text{where} \quad q_1 = \frac{f''}{f}, \tag{2.15}$$

and $v = \mathcal{I}(W)$ is a solution of the equation

$$\big(\square - q_2(x)\big)v = 0, \quad \text{where} \quad q_2 = \frac{(f^{-1})''}{f^{-1}} = 2\Big(\frac{f'}{f}\Big)^2 - q_1. \tag{2.16}$$

Theorem 2.2 [**25**]. *Let u be a scalar solution of the Klein-Gordon equation* (2.15) *in a simply connected domain Ω. Then a scalar solution v of the associated Klein-Gordon equation* (2.16) *such that $u + \mathbf{j}v$ is a solution of* (2.5) *in Ω, can be constructed according to the formula*

$$v = -f^{-1}\overline{A}_h\big[\mathbf{j}f^2\partial_{\bar z}(f^{-1}u)\big].$$

It is unique up to an additive term cf^{-1} where c is an arbitrary complex constant.

Vice versa, given a solution v of (2.16)*, the corresponding solution u of* (2.15) *such that $u + \mathbf{j}v$ is a solution of* (2.5)*, has the form*

$$u = -f\overline{A}_h\big[\mathbf{j}f^{-2}\partial_{\bar z}(fv)\big],$$

up to an additive term cf.

2.3. *Integration and formal powers*

Let (F,G) be a generating pair. Its adjoint generating pair $(F,G)^* = (F^*,G^*)$ is defined by the formulae

$$F^* = -\frac{2\overline{F}}{F\overline{G} - \overline{F}G}, \qquad G^* = \frac{2\overline{G}}{F\overline{G} - \overline{F}G}.$$

For our special case of generating pairs we have $(f, \mathbf{j}/f)^* = (f\mathbf{j}, 1/f)$ and $(1/f, f\mathbf{j})^* = (\mathbf{j}/f, f)$.

The (F,G)-integral is defined as

$$\int_\Gamma W d_{(F,G)}z = \frac{1}{2}\left(F(z_1)\,\mathcal{R}\left(\int_\Gamma G^*W\,dz\right) + G(z_1)\,\mathcal{R}\left(\int_\Gamma F^*W\,dz\right)\right)$$

where Γ is a rectifiable curve leading from z_0 to z_1.

If $W = \varphi F + \psi G$ is a bicomplex (F,G)-pseudoanalytic function where φ and ψ are scalar functions then

$$\int_{z_0}^{z} \overset{\circ}{W}\,d_{(F,G)}z = W(z) - \varphi(z_0)F(z) - \psi(z_0)G(z),$$

and this integral is path-independent and represents the (F,G)-antiderivative of $\overset{\circ}{W}$.

Definition 2.3 [**2**]. The formal power $Z_m^{(0)}(a,z_0;z)$ with center at z_0, coefficient a and exponent 0 is defined as the linear combination of the generators F_m, G_m with scalar constant coefficients λ, μ chosen so that $\lambda F_m(z_0)+\mu G_m(z_0)=a$. The formal powers with exponents $n=0,1,2,\dots$ are defined by the recursion formula

$$Z_m^{(n+1)}(a,z_0;z)=(n+1)\int_{z_0}^{z} Z_{m+1}^{(n)}(a,z_0;\zeta)\,d_{(F_m,G_m)}\zeta.$$

This definition implies the following properties.

(1) $Z_m^{(n)}(a,z_0;z)$ is an (F_m,G_m)-bicomplex pseudoanalytic function of z, i.e., it satisfies the hyperbolic Vekua equation determined by (F_m,G_m) (see [**23**]).

(2) If a' and a'' are scalar constants, then

$$Z_m^{(n)}(a'+\mathbf{j}a'',z_0;z)=a'Z_m^{(n)}(1,z_0;z)+a''Z_m^{(n)}(\mathbf{j},z_0;z).$$

(3) The formal powers satisfy the differential relations

$$\frac{d_{(F_m,G_m)}Z_m^{(n)}(a,z_0;z)}{dz}=nZ_{m+1}^{(n-1)}(a,z_0;z).$$

(4) The asymptotic formulae

$$Z_m^{(n)}(a,z_0;z)\sim a(z-z_0)^n,\quad z\to z_0$$

hold.

In the following we denote $Z_0^{(n)}(a,z_0;z)$ by $Z^{(n)}(a,z_0;z)$.

Since we are primarily interested in the hyperbolic Vekua equation (2.5), observe that as was explained above, a corresponding generating sequence is periodic having the form $(F_n,G_n)=(f,\mathbf{j}/f)$ when n is even, and $(F_n,G_n)=(1/f,f\mathbf{j})$ when n is odd. Using the construction from [**2**] given there for an analogous elliptic situation (see also [**23**, Sect. 4.2]) we obtain that the corresponding formal powers admit the following elegant representation. We consider the formal powers with the center in the origin and for simplicity assume that $f(0)=1$. Define the following systems of functions

$$X^{(0)}(x)\equiv\widetilde{X}^{(0)}(x)\equiv 1, \tag{2.17}$$

$$X^{(n)}(x)=n\int_0^x X^{(n-1)}(s)\left(f^2(s)\right)^{(-1)^n}\,\mathrm{d}s, \tag{2.18}$$

$$\widetilde{X}^{(n)}(x)=n\int_0^x \widetilde{X}^{(n-1)}(s)\left(f^2(s)\right)^{(-1)^{n-1}}\,\mathrm{d}s. \tag{2.19}$$

Then the formal powers corresponding to (2.5) have the following form. For $\alpha=\alpha'+\mathbf{j}\alpha''$ we have

$$Z^{(n)}(\alpha,0;z)=f(x)\mathcal{R}\big({}_*Z^{(n)}(\alpha,0;z)\big)+\frac{\mathbf{j}}{f(x)}\mathcal{I}\big({}_*Z^{(n)}(\alpha,0;z)\big), \tag{2.20}$$

where

$${}_*Z^{(n)}(\alpha,0;z)=\alpha'\sum_{k=0}^{n}\binom{n}{k}X^{(n-k)}\mathbf{j}^k t^k+\mathbf{j}\alpha''\sum_{k=0}^{n}\binom{n}{k}\widetilde{X}^{(n-k)}\mathbf{j}^k t^k \qquad \text{for an odd } n \tag{2.21}$$

and

$${}_*Z^{(n)}(\alpha,0;z)=\alpha'\sum_{k=0}^{n}\binom{n}{k}\widetilde{X}^{(n-k)}\mathbf{j}^k t^k+\mathbf{j}\alpha''\sum_{k=0}^{n}\binom{n}{k}X^{(n-k)}\mathbf{j}^k t^k \qquad \text{for an even } n. \tag{2.22}$$

For any $\alpha\in\mathbb{B}$ and $n\in\mathbb{N}_0=\mathbb{N}\cup\{0\}$ the formal power (2.20) is a solution of (2.5).

Remark 2.4. Formulae (2.20)–(2.22) clearly generalize the binomial representation for the analytic powers αz^n. If one chooses $f \equiv 1$ then $Z^{(n)}(\alpha, 0; z) = \alpha z^n$.

The formal powers $Z^{(n)}(\alpha, 0; z)$ can be written with the use of generalized wave polynomials [**18**]. Consider two families of functions $\{\varphi_k\}_{k=0}^{\infty}$ and $\{\psi_k\}_{k=0}^{\infty}$ constructed from (2.17), (2.18) and (2.19) by

$$\varphi_k(x) = \begin{cases} f(x) X^{(k)}(x), & k \text{ odd}, \\ f(x) \widetilde{X}^{(k)}(x), & k \text{ even}, \end{cases} \tag{2.23}$$

and

$$\psi_k(x) = \begin{cases} \dfrac{\widetilde{X}^{(k)}(x)}{f(x)}, & k \text{ odd}, \\ \dfrac{X^{(k)}(x)}{f(x)}, & k \text{ even}. \end{cases} \tag{2.24}$$

The following functions

$$u_0 = \varphi_0(x), \quad u_{2m-1}(x,t) = \sum_{\text{even } k=0}^{m} \binom{m}{k} \varphi_{m-k}(x) t^k, \quad u_{2m}(x,t) = \sum_{\text{odd } k=1}^{m} \binom{m}{k} \varphi_{m-k}(x) t^k, \tag{2.25}$$

and

$$v_0 = \psi_0(x), \quad v_{2m-1}(x,t) = \sum_{\text{even } k=0}^{m} \binom{m}{k} \psi_{m-k}(x) t^k, \quad v_{2m}(x,t) = \sum_{\text{odd } k=1}^{m} \binom{m}{k} \psi_{m-k}(x) t^k, \tag{2.26}$$

are called generalized wave polynomials (the wave polynomials are introduced below, in Remark 2.9). It is easy to see that

$$Z^{(0)}(\alpha, 0; z) = \alpha' u_0(x,t) + \mathbf{j}\alpha'' v_0(x,t), \tag{2.27}$$

$$Z^{(n)}(\alpha, 0; z) = \alpha' u_{2n-1}(x,t) + \alpha'' u_{2n}(x,t) + \mathbf{j}\big(\alpha' v_{2n}(x,t) + \alpha'' v_{2n-1}(x,t)\big), \quad n \geq 1. \tag{2.28}$$

The following parity relations hold for generalized wave polynomials.

$$u_0(x,-t) = u_0(x,t), \qquad u_{2n-1}(x,-t) = u_{2n-1}(x,t), \qquad u_{2n}(x,-t) = -u_{2n}(x,t). \tag{2.29}$$

It is worth to mention the following result obtained in [**22**] (for additional details and simpler proof see [**23**] and [**24**]) which establishes the relation of the systems of functions $\{\varphi_k\}_{k=0}^{\infty}$ and $\{\psi_k\}_{k=0}^{\infty}$ to the Sturm-Liouville equation.

Theorem 2.5 [**22**]. *Let q be a continuous complex valued function of an independent real variable $x \in [a,b]$ and λ be an arbitrary complex number. Suppose there exists a solution f of the equation*

$$f'' - qf = 0 \tag{2.30}$$

on (a,b) such that $f \in C^2(a,b) \cap C^1[a,b]$ and $f(x) \neq 0$ for any $x \in [a,b]$. Then the general solution $g \in C^2(a,b) \cap C^1[a,b]$ of the equation

$$g'' - qg = \lambda g$$

on (a,b) has the form

$$g = c_1 g_1 + c_2 g_2$$

where c_1 *and* c_2 *are arbitrary complex constants,*

$$g_1 = \sum_{k=0}^{\infty} \frac{\lambda^k}{(2k)!} \varphi_{2k} \qquad \text{and} \qquad g_2 = \sum_{k=0}^{\infty} \frac{\lambda^k}{(2k+1)!} \varphi_{2k+1} \tag{2.31}$$

and both series converge uniformly on $[a, b]$ *together with the series of the first derivatives which have the form*

$$g_1' = f' + \sum_{k=1}^{\infty} \frac{\lambda^k}{(2k)!} \left(\frac{f'}{f} \varphi_{2k} + 2k\, \psi_{2k-1} \right) \qquad \text{and}$$

$$g_2' = \sum_{k=0}^{\infty} \frac{\lambda^k}{(2k+1)!} \left(\frac{f'}{f} \varphi_{2k+1} + (2k+1)\, \psi_{2k} \right). \tag{2.32}$$

The series of the second derivatives converge uniformly on any segment $[a_1, b_1] \subset (a, b)$.

Representations (2.31) and (2.32), also known as the SPPS method, present an efficient and highly competitive technique for solving a variety of spectral and scattering problems related to Sturm-Liouville equations, see [**9**], [**14**], [**17**], [**19**], [**24**] and [**27**] and references therein.

2.4. *The Goursat problem for the hyperbolic Cauchy-Riemann system and a Runge type theorem*

For the construction of the kernels of the transmutation operators we are interested in solving corresponding Goursat problems. Our results concerning the Goursat problems for equation (2.5) arising as a natural step in that construction are based on the following simplest case.

Consider the hyperbolic Cauchy-Riemann system

$$\partial_{\bar{z}} w = 0 \quad \Longleftrightarrow \quad \begin{cases} u_x = v_t \\ u_t = v_x \end{cases} \tag{2.33}$$

studied in a considerable number of publications (see, e.g., [**31**], [**36**], [**47**]). It is easy to see that its general solution can be written as follows

$$\begin{aligned} w(x,t) &= \frac{1}{2} \left(\Phi\left(\frac{x+t}{2}\right) + \Psi\left(\frac{x-t}{2}\right) + \mathbf{j}\left(\Phi\left(\frac{x+t}{2}\right) - \Psi\left(\frac{x-t}{2}\right)\right)\right) \\ &= P^+ \Phi\left(\frac{x+t}{2}\right) + P^- \Psi\left(\frac{x-t}{2}\right) \end{aligned} \tag{2.34}$$

where Φ and Ψ are arbitrary continuously differentiable scalar functions.

Considering the values of w on the characteristics $x = t$ and $x = -t$ we see that

$$w(x, x) = P^+ \Phi(x) + P^- \Psi(0) \quad \text{and} \quad w(x, -x) = P^+ \Phi(0) + P^- \Psi(x). \tag{2.35}$$

Thus, we arrive at the following correct statement of the Goursat problem. *Find a solution* w *of* (2.33) *in a closed square* $\overline{R}$ *with the vertices* $(\pm 2b, 0)$ *and* $(0, \pm 2b)$, $b > 0$ *satisfying the conditions* (2.35) *where* Φ *and* Ψ *are arbitrary scalar functions such that* $\Phi, \Psi \in C^1[-b, b]$.

PROPOSITION 2.6. *The unique solution of the Goursat problem* (2.33), (2.35) *has the form* (2.34).

Proof. The fact that w is a solution of (2.33) follows from (2.34). The conditions (2.35) are obviously satisfied. To prove the uniqueness one considers the Goursat problem with the homogeneous conditions and finds out that Φ and Ψ from the representation (2.34) of the general solution of (2.33) satisfy the equations $P^{+}\Phi(x)+P^{-}\Psi(0)\equiv 0$ and $P^{+}\Phi(0)+P^{-}\Psi(x)\equiv 0$, $x\in[-b,b]$ from which both Φ and Ψ are necessarily trivial. □

This proposition is not entirely new. As a kind of analogue of the Cauchy integral formula for analytic functions it was discussed, e.g., in [**36**] in the case $w:\mathbb{D}\to\mathbb{D}$. For our purposes it is more convenient to consider this simple representation of any hyperbolic analytic function via its values on the characteristics from the point of view of the Goursat problem.

The following result is a direct corollary of Proposition 2.6.

PROPOSITION 2.7. *Let Φ and Ψ be scalar functions defined on $[-b,b]$ and admitting there the uniformly convergent power series expansions*

$$\Phi(x)=\sum_{n=0}^{\infty}\alpha_n x^n \quad \text{and} \quad \Psi(x)=\sum_{n=0}^{\infty}\beta_n x^n.$$

Then the unique solution of the Goursat problem (2.33), (2.35) *has the form*

$$w=\sum_{n=0}^{\infty}a_n z^n, \tag{2.36}$$

where

$$a_n=\frac{1}{2^n}\left(P^{+}\alpha_n+P^{-}\beta_n\right). \tag{2.37}$$

Proof. Due to Proposition 2.6 we have

$$\begin{aligned} w(x,t)&=P^{+}\sum_{n=0}^{\infty}\alpha_n\left(\frac{x+t}{2}\right)^n+P^{-}\sum_{n=0}^{\infty}\beta_n\left(\frac{x-t}{2}\right)^n\\ &=\sum_{n=0}^{\infty}\left(\frac{\alpha_n}{2^n}P^{+}(x+t)^n+\frac{\beta_n}{2^n}P^{-}(x-t)^n\right). \end{aligned}$$

Notice that $P^{\pm}z=P^{\pm}(x\pm t)$ and hence $P^{\pm}z^n=P^{\pm}(x\pm t)^n$. Thus,

$$w=\sum_{n=0}^{\infty}\left(\frac{\alpha_n}{2^n}P^{+}z^n+\frac{\beta_n}{2^n}P^{-}z^n\right)=\sum_{n=0}^{\infty}\frac{1}{2^n}\left(P^{+}\alpha_n+P^{-}\beta_n\right)z^n.$$

□

REMARK 2.8. Under the conditions of Proposition 2.7 we have that the unique solution w of the Goursat problem (2.33), (2.35) can be represented as a uniformly convergent in $\overline{R}$ Taylor series (2.36) where

$$a_n=\frac{w^{[n]}(0)}{n!}. \tag{2.38}$$

Indeed, let us prove this formula for the coefficients. Due to (2.12) we have that $w^{[n]}=\mathbf{j}^n\partial_t^n w$ and due to (2.34),

$$\mathbf{j}^n\partial_t^n w=\frac{1}{2^n}\left(P^{+}\mathbf{j}^n\Phi^{(n)}\left(\frac{x+t}{2}\right)+P^{-}\mathbf{j}^n(-1)^n\Psi^{(n)}\left(\frac{x-t}{2}\right)\right).$$

Now we notice that $P^{+}\mathbf{j}^n=P^{+}$ and $P^{-}\mathbf{j}^n(-1)^n=P^{-}$ for any $n=0,1,\ldots$. Thus, $w^{[n]}(0)=\frac{n!}{2^n}\left(P^{+}\alpha_n+P^{-}\beta_n\right)=n!a_n$.

REMARK 2.9. It is possible to give another representation for the solution of the Goursat problem, namely, a representation as a series in wave polynomials, cf. [**18**, Proposition 1]. Recall that the following polynomials

$$p_0(x,t)=1, \quad p_{2m-1}(x,t)=\mathcal{R}\big((x+\mathbf{j}t)^m\big), \quad p_{2m}(x,t)=\mathcal{I}\big((x+\mathbf{j}t)^m\big),\ m\geq 1,$$

are called wave polynomials. They may be also written as

$$p_0(x,t)=1, \quad p_{2m-1}(x,t)=\sum_{\text{even } k=0}^{m}\binom{m}{k}x^{m-k}t^k, \quad p_{2m}(x,t)=\sum_{\text{odd } k=1}^{m}\binom{m}{k}x^{m-k}t^k, \quad (2.39)$$

cf. with the definition of generalized wave polynomials (2.25), (2.26). Since $z^n=p_{2n-1}(x,t)+\mathbf{j}p_{2n}(x,t)$ for $n\geq 1$, we obtain from (2.37) that

$$w(x,t)=(P^+\alpha_0+P^-\beta_0)p_0(x,t) \\ +\sum_{n=1}^{\infty}\frac{P^+\alpha_n+P^-\beta_n}{2^n}p_{2n-1}(x,t)+\sum_{n=1}^{\infty}\frac{P^+\alpha_n-P^-\beta_n}{2^n}p_{2n}(x,t).$$

Obviously, an arbitrary continuously differentiable solution of (2.33) cannot be represented globally as a uniformly convergent power series of the form (2.36). Nevertheless the following analogue of the Runge approximation theorem from classical complex analysis is a corollary of the preceding proposition.

PROPOSITION 2.10. *Let $w\in C^1(\overline{R})$ be a solution of* (2.33) *in R. Then there exists a sequence of polynomials $P_N=\sum_{n=0}^{N}a_nz^n$ uniformly convergent to w in $\overline{R}$.*

Proof. We need to prove that for any $\varepsilon>0$ there exist such a number N and coefficients a_n, $n=0,1,\dots,N$ that $|w(x,t)-P_N(x,t)|<\varepsilon$ for any point $(x,t)\in\overline{R}$. Let the scalar functions Φ and Ψ defined on $[-b,b]$ be such that (2.35) hold. As was shown above such scalar functions always exist and under the condition of the proposition they are obviously from $C^1[-b,b]$. We choose $\varepsilon>0$ and such $\varepsilon_{1,2}>0$ that $\varepsilon=(\varepsilon_1+\varepsilon_2)/2$. According to the Weierstrass theorem there exists such a number N and such polynomials p_1 and p_2 of order not greater than a certain N that $|\Phi(x)-p_1(x)|_{\mathbb{C}}<\varepsilon_1$ and $|\Psi(x)-p_2(x)|_{\mathbb{C}}<\varepsilon_2$, $-b\leq x\leq b$. Due to Proposition 2.7 the unique solution $\widetilde{w}$ of the Goursat problem for equation (2.33) with the boundary conditions

$$\widetilde{w}(x,x)=P^+p_1(x)+P^-p_2(0), \qquad \widetilde{w}(x,-x)=P^+p_1(0)+P^-p_2(x)$$

has the form $\widetilde{w}=P_N$ where $P_N=\sum_{n=0}^{N}\frac{1}{2^n}\left(P^+\alpha_n+P^-\beta_n\right)z^n$ with α_n and β_n being the coefficients of the polynomials p_1 and p_2 respectively.

Consider

$$\begin{aligned}|w(x,t)-\widetilde{w}(x,t)|&=|w(x,t)-P_N(x,t)|\\ &=\left|P^+\Phi\left(\frac{x+t}{2}\right)+P^-\Psi\left(\frac{x-t}{2}\right)-P^+P_N^+(x,t)-P^-P_N^-(x,t)\right|\\ &=\frac{1}{2}\left(\left|\Phi\left(\frac{x+t}{2}\right)-P_N^+(x,t)\right|_{\mathbb{C}}+\left|\Psi\left(\frac{x-t}{2}\right)-P_N^-(x,t)\right|_{\mathbb{C}}\right),\end{aligned}$$

where we have used (2.1) and (2.2). Notice that $P_N^+(x,t)=\sum_{n=0}^{N}\alpha_n\left(\frac{x+t}{2}\right)^n=p_1\left(\frac{x+t}{2}\right)$ and $P_N^-(x,t)=\sum_{n=0}^{N}\beta_n\left(\frac{x-t}{2}\right)^n=p_2\left(\frac{x-t}{2}\right)$. Thus, $|w(x,t)-P_N(x,t)|<\frac{1}{2}(\varepsilon_1+\varepsilon_2)=\varepsilon$. □

3. *Transmutation operators*

We give a definition of a transmutation operator from [**27**] which is a modification of the definition given by Levitan [**32**] adapted to the purposes of the present work. Let E be a linear topological space and E_1 its linear subspace (not necessarily closed). Let A and B be linear operators: $E_1 \to E$.

Definition 3.1. A linear invertible operator T defined on the whole E such that E_1 is invariant under the action of T is called a transmutation operator for the pair of operators A and B if it fulfills the following two conditions.

(i) Both the operator T and its inverse T^{-1} are continuous in E;

(ii) The following operator equality is valid

$$AT = TB \tag{3.1}$$

or which is the same

$$A = TBT^{-1}.$$

Our main interest concerns the situation when $A = -\frac{d^2}{dx^2} + q(x)$, $B = -\frac{d^2}{dx^2}$, and q is a continuous complex-valued function. Hence for our purposes it will be sufficient to consider the functional space $E = C[a,b]$ with the topology of uniform convergence and its subspace E_1 consisting of functions from $C^2\,[a,b]$. One of the possibilities to introduce a transmutation operator on E was considered by Lions [**33**] and later on in other references (see, e.g., [**34**]), and consists in constructing a Volterra integral operator corresponding to a midpoint of the segment of interest. As we begin with this transmutation operator it is convenient to consider a symmetric segment $[-b,b]$ and hence the functional space $E = C[-b,b]$. It is worth mentioning that other well known ways to construct the transmutation operators (see, e.g., [**32**], [**44**]) imply imposing initial conditions on the functions and consequently lead to transmutation operators satisfying (3.1) only on subclasses of E_1.

Thus, consider the space $E = C[-b,b]$. In [**6**] and [**26**] a parametrized family of transmutation operators for the defined above A and B was studied. Operators of this family can be realized in the form of the Volterra integral operator

$$\mathbf{T}_h u(x) = u(x) + \int_{-x}^{x} \mathbf{K}(x,t;h)u(t)dt \tag{3.2}$$

where $\mathbf{K}(x,t;h) = \mathbf{H}\big(\frac{x+t}{2}, \frac{x-t}{2}; h\big)$, h is a complex parameter, $|t| \le |x| \le b$ and $\mathbf{H}$ is the unique solution of the Goursat problem

$$\frac{\partial^2 \mathbf{H}(u,v;h)}{\partial u\, \partial v} = q(u+v)\mathbf{H}(u,v;h), \tag{3.3}$$

$$\mathbf{H}(u,0;h) = \frac{h}{2} + \frac{1}{2}\int_0^u q(s)\,ds, \qquad \mathbf{H}(0,v;h) = \frac{h}{2}. \tag{3.4}$$

If the potential q is continuously differentiable, the kernel $\mathbf{K}$ itself is a solution of the Goursat problem

$$\left(\frac{\partial^2}{\partial x^2} - q(x)\right)\mathbf{K}(x,t;h) = \frac{\partial^2}{\partial t^2}\mathbf{K}(x,t;h), \tag{3.5}$$

$$\mathbf{K}(x,x;h) = \frac{h}{2} + \frac{1}{2}\int_0^x q(s)\,ds, \qquad \mathbf{K}(x,-x;h) = \frac{h}{2}. \tag{3.6}$$

If the potential q is n times continuously differentiable, the kernel $\mathbf{K}(x,t;h)$ is $n+1$ times continuously differentiable with respect to both independent variables.

REMARK 3.2. In the case $h = 0$ the operator $\mathbf{T}_h$ coincides with the transmutation operator studied in [**34**, Chap. 1, Sect. 2]. In [**32**], [**33**], [**44**] it was established that in the case $q \in C^1[-b, b]$ the Volterra-type integral operator (3.2) is a transmutation in the sense of Definition 3.1 on the space $C^2[-b, b]$ if and only if the integral kernel $\mathbf{K}(x, t)$ satisfies the Goursat problem (3.5), (3.6).

The following proposition shows that it is sufficient to know the transmutation operator $\mathbf{T}_{h_1}$ or its integral kernel $\mathbf{K}_{h_1}$ for some particular parameter h_1 to be able to construct transmutation operators $\mathbf{T}_h$ or their integral kernels $\mathbf{K}_h$ for arbitrary values of the parameter h.

PROPOSITION 3.3 [**6**], [**27**]. *The operators $\mathbf{T}_{h_1}$ and $\mathbf{T}_{h_2}$ are related by the expression*

$$\mathbf{T}_{h_2} u = \mathbf{T}_{h_1}\left[u(x) + \frac{h_2 - h_1}{2}\int_{-x}^{x} u(t)\,dt\right]$$

valid for any $u \in C[-b, b]$.

The integral kernels $\mathbf{K}(x, t; h_1)$ and $\mathbf{K}(x, t; h_2)$ are related by the expression

$$\mathbf{K}(x, t; h_2) = \frac{h_2 - h_1}{2} + \mathbf{K}(x, t; h_1) + \frac{h_2 - h_1}{2}\int_{t}^{x}\big(\mathbf{K}(x, s; h_1) - \mathbf{K}(x, -s; h_1)\big)\,ds.$$

The following theorem states that the operators $\mathbf{T}_h$ are indeed transmutations in the sense of Definition 3.1.

THEOREM 3.4. *Let $q \in C[-b, b]$. Then the operator $\mathbf{T}_h$ given by* (3.2) *satisfies the equality*

$$\left(-\frac{d^2}{dx^2} + q(x)\right)\mathbf{T}_h[u] = \mathbf{T}_h\left[-\frac{d^2}{dx^2}(u)\right] \tag{3.7}$$

for any $u \in C^2[-b, b]$.

REMARK 3.5. This theorem under additional assumptions on q was proved in [**6**] and [**26**]. In [**27**] it was shown that (3.7) holds for any h whenever it holds for some particular value h_1.

Proof. It follows from Proposition 3.3 that it is sufficient to prove (3.7) for one particular h, see [**27**, Proof of Theorem 5.6] for details. We take $h = 0$ and the operator $\mathbf{T} := \mathbf{T}_0$. Let $\mathbf{K}$ denotes the integral kernel of $\mathbf{T}$. There exists a sequence of potentials $\{q_n\}_{n\in\mathbb{N}} \subset C^1[-b, b]$ such that $q_n \to q$, $n \to \infty$ uniformly on $[-b, b]$. Let $\mathbf{K}_{q_n}$ and $\mathbf{T}_{q_n}$ be the integral kernels and the transmutation operators corresponding to the potentials q_n. Then $\mathbf{K}_{q_n} \to \mathbf{K}$ uniformly and $\mathbf{T}_n \to \mathbf{T}$ in the operator norm, see, e.g., [**34**, Chap. 1, Sect. 2]. Take a function $u \in C^2[-b, b]$. For the operators $\mathbf{T}_{q_n}$ the following equality holds [**26**, Theorem 6]

$$\left(-\frac{d^2}{dx^2} + q_n(x)\right)\mathbf{T}_{q_n}[u] = \mathbf{T}_{q_n}\left[-\frac{d^2}{dx^2}(u)\right],$$

which with the use of (3.2) may be rewritten as

$$\frac{d^2}{dx^2}\left(\int_{-x}^{x}\mathbf{K}_{q_n}(x, t)u(t)\,dt\right) = -u''(x) + q_n(x)\mathbf{T}_{q_n}[u](x) + \mathbf{T}_{q_n}[u''](x).$$

Let $y_n(x) := \int_{-x}^{x}\mathbf{K}_{q_n}(x, t)u(t)\,dt$ and $z_n := -u'' + q_n\mathbf{T}_{q_n}[u] + \mathbf{T}_{q_n}[u'']$. Then $y_n \to y := \int_{-x}^{x}\mathbf{K}(x, t)u(t)\,dt$ and $z_n \to z := -u'' + q\mathbf{T}[u] + \mathbf{T}[u'']$, $n \to \infty$. Moreover, functions y_n satisfy the initial conditions $y_n(0) = y_n'(0) = 0$. Now (3.7) follows from the fact that the operator ∂_x^2 with the domain $\{y \in C^2[-b,b] : y(0) = y'(0) = 0\}$ is closed, see, e.g., [**16**]. □

The following theorem summarizes the mapping properties of the operators $\mathbf{T}_h$ establishing that there exists a value of the parameter h such that $\mathbf{T}_h$ maps x^k to φ_k defined by (2.23).

Theorem 3.6 [**6**], [**26**]. *Let q be a continuous complex valued function of an independent real variable $x \in [-b,b]$ for which there exists a particular solution f of* (2.30) *such that $f \in C^2[-b,b]$, $f \neq 0$ on $[-b,b]$ and normalized as $f(0) = 1$. Denote $h := f'(0) \in \mathbb{C}$. Suppose $\mathbf{T}_h$ is the operator defined by* (3.2) *and φ_k, $k \in \mathbb{N}_0$ are functions defined by* (2.23). *Then*

$$\mathbf{T}_h x^k = \varphi_k(x) \qquad \text{for any } k \in \mathbb{N}_0. \tag{3.8}$$

Moreover, $\mathbf{T}_h$ maps a solution v of an equation $v'' + \omega^2 v = 0$, where ω is a complex number, into a solution u of the equation $u'' - q(x)u + \omega^2 u = 0$ with the following correspondence of the initial values

$$u(0) = v(0), \qquad u'(0) = v'(0) + hv(0).$$

Remark 3.7. The mapping property (3.8) of the transmutation operator allows one to see that the SPPS representations (2.31) from Theorem 2.5 are nothing but the images of Taylor expansions of the functions $\cosh\sqrt{\lambda}x$ and $\frac{1}{\sqrt{\lambda}}\sinh\sqrt{\lambda}x$. Moreover, equality (3.8) is behind a new method for solving Sturm-Liouville problems proposed in [**28**].

In what follows we assume that $f \in C^2[-b,b]$, $f \neq 0$ on $[-b,b]$, $f(0) = 1$ and denote $h := f'(0) \in \mathbb{C}$. Any such function is associated with an operator $\mathbf{T}_h$. For convenience, from now on we will write T_f instead of $\mathbf{T}_h$ and the integral kernel of T_f will be denoted by $\mathbf{K}_f$. Together with the function f let us consider the function $1/f$. Notice that if f is a solution of (2.30) with the potential $q_f = f''/f$, the function $1/f$ is a solution of the equation

$$\left(-\frac{d^2}{dx^2} + q_{1/f}(x)\right) u = 0$$

with the Darboux associated potential

$$q_{1/f} = -q_f + 2\left(\frac{f'}{f}\right)^2.$$

Consider the transmutation operator $T_{1/f}$ which satisfies the equality

$$\left(-\frac{d^2}{dx^2} + q_{1/f}(x)\right) T_{1/f}[u] = T_{1/f}\left[-\frac{d^2}{dx^2}(u)\right]$$

for any $u \in C^2[-b,b]$ and transforms x^k into the functions $\psi_k(x)$, $k \in \mathbb{N}_0$ defined by (2.24), i.e.

$$T_{1/f} x^k = \psi_k(x) \qquad \text{for any } k \in \mathbb{N}_0. \tag{3.9}$$

It follows from (3.8), (3.9) and definitions (2.25), (2.26) and (2.39) that generalized wave polynomials are images of wave polynomials under the action of operators T_f and $T_{1/f}$, i.e.

$$T_f p_n = u_n \quad \text{and} \quad T_{1/f} p_n = v_n \quad \text{for any } n \geq 0. \tag{3.10}$$

In [**26**] explicit formulas were obtained for the kernel $\mathbf{K}_{1/f}(x,t)$ in terms of $\mathbf{K}_f(x,t)$. In order to obtain a simpler expression for the integral kernel $\mathbf{K}_{1/f}(x,t)$ we assumed that the original integral kernel $\mathbf{K}_f(x,t)$ is known in the larger domain than required by definition (3.2). Namely, suppose that the function $\mathbf{K}_f(x,t)$ is known and is continuously differentiable in the domain $\overline{\Pi}:\ -b \leq x \leq b, -b \leq t \leq b$. It is worth mentioning that such continuation of the integral kernel to the domain $\bar{\Pi}$ is always possible due to (3.3), (3.4) and the general theory of Goursat problems. We refer the reader to [**26**] for further details. In Corollary 7.2 based on pseudoanalytic function theory we obtain another expression for the integral kernel $\mathbf{K}_{1/f}(x,t)$ suitable in the case when the integral kernel $\mathbf{K}_f(x,t)$ is known only in the natural domain $|t| \le |x| \le b$.

PROPOSITION 3.8 [**26**]. *Under the conditions of Theorem 3.6 the transmutation operators* T_f *and* $T_{1/f}$ *are related by the expressions*

$$T_{1/f}[u](x) = \frac{1}{f(x)}\left(\int_0^x f(\eta)T_f[u'](\eta)\,d\eta + u(0)\right)$$

and

$$T_f[u](x) = f(x)\left(\int_0^x \frac{1}{f(\eta)}T_{1/f}[u'](\eta)\,d\eta + u(0)\right),$$

valid for any $u \in C^1[-b,b]$, *and their integral kernels* $\mathbf{K}_f(x,t) := \mathbf{K}_f(x,t;h)$ *and* $\mathbf{K}_{1/f}(x,t) := \mathbf{K}_{1/f}(x,t;-h)$ *are related by the expressions*

$$\mathbf{K}_{1/f}(x,t) = -\frac{1}{f(x)}\left(\int_{-t}^x \partial_t\mathbf{K}_f(s,t)f(s)\,ds + \frac{f'(0)}{2}f(-t)\right) \tag{3.11}$$

and

$$\mathbf{K}_f(x,t) = -f(x)\left(\int_{-t}^x \partial_t\mathbf{K}_{1/f}(s,t)\frac{1}{f(s)}\,ds - \frac{f'(0)}{2f(-t)}\right). \tag{3.12}$$

The following commutation relations immediately follow from Proposition 3.8.

COROLLARY 3.9 [**26**]. *The following operator equalities hold on* $C^1[-b,b]$:

$$\begin{aligned}\partial_x f T_{1/f} &= f T_f \partial_x\\ \partial_x \frac{1}{f} T_f &= \frac{1}{f} T_{1/f} \partial_x.\end{aligned}$$

EXAMPLE 3.10 [**26**]. Take $f(x) = x+1$ and some segment $[-b,b] \subset (-1,1)$. Then $q_f = 0$ and $q_{1/f} = \frac{2}{(x+1)^2}$. For the potential q_f the transmutation operator $\mathbf{T}_0$ is obviously the identity operator, hence $\mathbf{K}(x,t;0) = 0$ and from Proposition 3.3 we obtain

$$\mathbf{K}_f(x,t) := \mathbf{K}(x,t;1) = 1/2.$$

Hence we get from (3.11) that

$$\mathbf{K}_{1/f}(x,t) = \frac{t-1}{2(x+1)}.$$

It is easy to see that the integral kernels $\mathbf{K}_f$ and $\mathbf{K}_{1/f}$ satisfy the Goursat problems (3.5), (3.6) and relation (3.12) holds.

For more examples of transmutation integral kernels we refer the reader to [**26**].

4. *Goursat-to-Goursat transmutation operators*

Let $\widetilde{u}$ and u be solutions of the equations $\Box\widetilde{u} = 0$ and $(\Box - q(x))\,u = 0$ in $\overline{\mathbf{R}}$ where $\mathbf{R}$ is a square with the vertices $(\pm b, 0)$ and $(0, \pm b)$, respectively such that $u = T_f\widetilde{u}$. Consider the operator T_G mapping the Goursat data corresponding to $\widetilde{u}$ into the Goursat data corresponding to u,

$$T_G : \begin{pmatrix} \widetilde{u}(x,x) \\ \widetilde{u}(x,-x) \end{pmatrix} \longmapsto \begin{pmatrix} u(x,x) \\ u(x,-x) \end{pmatrix}.$$

Denote $\varphi(x) := \widetilde{u}(x,x)$, $\psi(x) := \widetilde{u}(x,-x)$, $\Phi(x) := u(x,x)$, $\Psi(x) := u(x,-x)$. We have $\varphi(0) = \psi(0)$ and $\Phi(0) = \Psi(0)$. The solution of the wave equation $\widetilde{u}$ is represented via its Goursat data as follows $\widetilde{u}(x,t) = \varphi(\frac{x+t}{2}) + \psi(\frac{x-t}{2}) - \varphi(0)$. Application of the operator T_f gives us the equalities

$$T_f\varphi\Big(\frac{x+t}{2}\Big) = \varphi\Big(\frac{x+t}{2}\Big) + \int_{-x}^{x} \mathbf{K}_f(x,\tau)\varphi\Big(\frac{\tau+t}{2}\Big)d\tau,$$

$$T_f\psi\Big(\frac{x-t}{2}\Big) = \psi\Big(\frac{x-t}{2}\Big) + \int_{-x}^{x} \mathbf{K}_f(x,\tau)\psi\Big(\frac{\tau-t}{2}\Big)d\tau$$

and $T_f[\varphi(0)] = \varphi(0)f$. Considering the value of these expressions when $t = x$ and $t = -x$ and introducing obvious changes of variables we obtain the following relations

$$\Phi(x) = \varphi(x) + 2\int_0^x \mathbf{K}_f(x,2t-x)\varphi(t)dt + \psi(0) + 2\int_{-x}^0 \mathbf{K}_f(x,2t+x)\psi(t)dt - \varphi(0)f(x)$$

and

$$\Psi(x) = \varphi(0) + 2\int_{-x}^0 \mathbf{K}_f(x,2t+x)\varphi(t)dt + \psi(x) + 2\int_0^x \mathbf{K}_f(x,2t-x)\psi(t)dt - \varphi(0)f(x).$$

Due to the boundedness of the operators T_f and T_f^{-1} as well as to the continuous dependence of the solutions of the Goursat problems under consideration on their respective Goursat data (see, e.g., [**46**, Sect. 15]) the operator T_G together with T_G^{-1} are bounded on vector functions from $C^1[-b,b] \times C^1[-b,b]$ equipped with a suitable, for example, maximum, norm.

It is easy to establish the following mapping properties of T_G,

$$T_G : \quad 2^{n-1}\begin{pmatrix} x^n \\ x^n \end{pmatrix} \longmapsto \begin{pmatrix} u_{2n-1}(x,x) \\ u_{2n-1}(x,x) \end{pmatrix} \tag{4.1}$$

and

$$T_G : \quad 2^{n-1}\begin{pmatrix} x^n \\ -x^n \end{pmatrix} \longmapsto \begin{pmatrix} u_{2n}(x,x) \\ -u_{2n}(x,x) \end{pmatrix}. \tag{4.2}$$

Indeed, consider the wave polynomial $p_{2n-1}(x,t)$, $n = 1,2,\ldots$, see (2.39). Its Goursat data have the form $p_{2n-1}(x,x) = p_{2n-1}(x,-x) = 2^{n-1}x^n$. The image of p_{2n-1} under the action of T_f is u_{2n-1}, see (3.10). Using the property (2.29) we obtain (4.1). Analogously, consideration of p_{2n} leads to (4.2).

PROPOSITION 4.1. *Let u be a regular solution of the equation $(\square - q(x))\,u = 0$ in $\overline{\mathbf{R}}$ such that its Goursat data admit the following series expansions*

$$\frac{1}{2}\left(u(x,x) + u(x,-x)\right) = c_0u_0(x,x) + \sum_{n=1}^{\infty} c_nu_{2n-1}(x,x), \tag{4.3}$$

and

$$\frac{1}{2}\left(u(x,x) - u(x,-x)\right) = \sum_{n=1}^{\infty} b_nu_{2n}(x,x), \tag{4.4}$$

both uniformly convergent on $[-b,b]$. Then for any $(x,t) \in \overline{\mathbf{R}}$,

$$u(x,t) = c_0u_0(x,t) + \sum_{n=1}^{\infty} \left(c_nu_{2n-1}(x,t) + b_nu_{2n}(x,t)\right) \tag{4.5}$$

and the series converges uniformly in $\overline{\mathbf{R}}$.

Proof. Notice that if (4.5) is valid then (4.3) and (4.4) hold. Indeed, due to (2.29)

$$\begin{aligned}\frac{1}{2}\left(u(x,x)+u(x,-x)\right) &= c_0u_0(x,x)\\ &+\frac{1}{2}\sum_{n=1}^{\infty}\left(c_n\left(u_{2n-1}(x,x)+u_{2n-1}(x,x)\right)+b_n\left(u_{2n}(x,x)-u_{2n}(x,x)\right)\right)\\ &= c_0u_0(x,x)+\sum_{n=1}^{\infty}c_nu_{2n-1}(x,x)\end{aligned}$$

and similarly (4.4) can be verified.

Suppose that (4.3) and (4.4) are valid. Then we have

$$\begin{pmatrix}u(x,x)\\u(x,-x)\end{pmatrix}=c_0\begin{pmatrix}u_0(x,x)\\u_0(x,x)\end{pmatrix}+\sum_{n=1}^{\infty}\left(c_n\begin{pmatrix}u_{2n-1}(x,x)\\u_{2n-1}(x,x)\end{pmatrix}+b_n\begin{pmatrix}u_{2n}(x,x)\\-u_{2n}(x,x)\end{pmatrix}\right).$$

From (4.1) and (4.2) we obtain

$$\begin{aligned}\begin{pmatrix}\varphi(x)\\\psi(x)\end{pmatrix} &:= T_G^{-1}\begin{pmatrix}u(x,x)\\u(x,-x)\end{pmatrix}=c_0\begin{pmatrix}1\\1\end{pmatrix}+\sum_{n=1}^{\infty}2^{n-1}\left(c_n\begin{pmatrix}x^n\\x^n\end{pmatrix}+b_n\begin{pmatrix}x^n\\-x^n\end{pmatrix}\right)\\ &=\begin{pmatrix}c_0+\sum_{n=1}^{\infty}2^{n-1}\left(c_n+b_n\right)x^n\\c_0+\sum_{n=1}^{\infty}2^{n-1}\left(c_n-b_n\right)x^n\end{pmatrix}.\end{aligned}\tag{4.6}$$

Under the conditions of the proposition and due to the boundedness of T_G^{-1} we have that both series in the last equality are uniformly convergent on $[-b,b]$. Then applying Proposition 1 from [**18**] we obtain that the unique solution $\widetilde{u}$ of the Goursat problem for the wave equation in $\overline{\mathbf{R}}$ and with the Goursat data $\binom{\varphi(x)}{\psi(x)}$ has the form

$$\widetilde{u}(x,t)=c_0p_0(x,t)+\sum_{n=1}^{\infty}\left(c_np_{2n-1}(x,t)+b_np_{2n}(x,t)\right)\tag{4.7}$$

and the series converges uniformly in $\overline{\mathbf{R}}$. Since $u=T_f\widetilde{u}$, $T_fp_n=u_n$ and T_f is bounded we obtain the uniform convergence of the series (4.5). □

5. *Transmutations for the Vekua operator*

By analogy with [**5**] we introduce the following pair of operators

$$\mathbf{V}_1=T_f\mathcal{R}+\mathbf{j}T_{1/f}\mathcal{I}\tag{5.1}$$

and

$$\mathbf{V}_2=T_{1/f}\mathcal{R}+\mathbf{j}T_f\mathcal{I}.\tag{5.2}$$

Proposition 5.1. *The following equalities hold for any* $\mathbb{B}$*-valued, continuously differentiable function* w *defined on* $\overline{\mathbf{R}}$.

$$\left(\partial_{\overline{z}}-\frac{\partial_{\overline{z}}f}{f}C\right)\mathbf{V}_1w=\mathbf{V}_2\left(\partial_{\overline{z}}w\right),\qquad\left(\partial_{\overline{z}}+\frac{\partial_zf}{f}C\right)\mathbf{V}_2w=\mathbf{V}_1\left(\partial_{\overline{z}}w\right).$$

$$\left(\partial_z-\frac{\partial_zf}{f}C\right)\mathbf{V}_1w=\mathbf{V}_2\left(\partial_zw\right),\qquad\left(\partial_z+\frac{\partial_{\overline{z}}f}{f}C\right)\mathbf{V}_2w=\mathbf{V}_1\left(\partial_zw\right).$$

Proof. The proof consists in a direct calculation with the aid of the commutation relations from Corollary 3.9. For example, for $w = u + \mathbf{j}v$ we have

$$\begin{aligned}\left(\partial_{\overline{z}} - \frac{\partial_{\overline{z}} f}{f} C\right) \mathbf{V}_1 w &= \frac{1}{2}\left(f\partial_{\overline{z}}\left(\frac{1}{f}\mathcal{R}\left(\mathbf{V}_1 w\right)\right) + \mathbf{j}\frac{1}{f}\partial_{\overline{z}}\left(f\mathcal{I}\left(\mathbf{V}_1 w\right)\right)\right)\\ &= \frac{1}{2}\left(f\partial_x\left(\frac{1}{f}T_f u\right) - \mathbf{j}\partial_t T_f u + \mathbf{j}\frac{1}{f}\partial_x\left(fT_{1/f}v\right) - \partial_t T_{1/f} v\right)\\ &= \frac{1}{2}\left(T_{1/f}\partial_x u - \mathbf{j}T_f\partial_t u + \mathbf{j}T_f\partial_x v - T_{1/f}\partial_t v\right) = \mathbf{V}_2\left(\partial_{\overline{z}} w\right).\end{aligned}$$

□

An immediate corollary from Proposition 5.1 is the fact that the operators $\mathbf{V}_1$ and $\mathbf{V}_2$ map the hyperbolic analytic functions satisfying (2.33) into the solutions of (2.4) and (2.10) respectively. Moreover, they map powers of the variable z into the corresponding formal powers.

PROPOSITION 5.2. *For any $z \in \overline{\mathbf{R}}$, $n \in \mathbb{N}_0$ and $a \in \mathbb{B}$ the following equalities are valid*

$$\mathbf{V}_1[az^n] = Z^{(n)}(a, 0; z) \quad \text{and} \quad \mathbf{V}_2[az^n] = Z_1^{(n)}(a, 0; z).$$

Proof. The proof consists in the observation that for $a = a' + \mathbf{j}b'$ and $z = x + \mathbf{j}t$ one has

$$az^n = (a' + \mathbf{j}b')\sum_{m=0}^{n}\binom{n}{m}x^{n-m}\mathbf{j}^m t^m$$

and the result follows from the formulas (2.20), (2.21), (2.22) by application of the mapping properties (3.8), (3.9). □

Notice that both operators $\mathbf{V}_1$ and $\mathbf{V}_2$ are bounded on the space of continuous functions. Indeed, consider

$$\begin{aligned}\left|\mathbf{V}_1 w\right| &= \frac{1}{2}\left(\left|(\mathbf{V}_1 w)^+\right|_{\mathbb{C}} + \left|(\mathbf{V}_1 w)^-\right|_{\mathbb{C}}\right)\\ &= \frac{1}{2}\left(\left|\mathcal{R}\left(\mathbf{V}_1 w\right) + \mathcal{I}\left(\mathbf{V}_1 w\right)\right|_{\mathbb{C}} + \left|\mathcal{R}\left(\mathbf{V}_1 w\right) - \mathcal{I}\left(\mathbf{V}_1 w\right)\right|_{\mathbb{C}}\right)\\ &\le \left|\mathcal{R}\left(\mathbf{V}_1 w\right)\right|_{\mathbb{C}} + \left|\mathcal{I}\left(\mathbf{V}_1 w\right)\right|_{\mathbb{C}} = \left|T_f u\right|_{\mathbb{C}} + \left|T_{1/f} v\right|_{\mathbb{C}}.\end{aligned}$$

From the boundedness of the operators T_f and $T_{1/f}$ we have

$$\max\left|\mathbf{V}_1 w\right| \le M\left(\max|u|_{\mathbb{C}} + \max|v|_{\mathbb{C}}\right)$$

where $M = \max\left\{\|T_f\|, \left\|T_{1/f}\right\|\right\}$. Since $|u|_{\mathbb{C}} \le \frac{1}{2}\left(|w^+|_{\mathbb{C}} + |w^-|_{\mathbb{C}}\right)$ and $|v|_{\mathbb{C}} \le \frac{1}{2}\left(|w^+|_{\mathbb{C}} + |w^-|_{\mathbb{C}}\right)$, we obtain $\max\left|\mathbf{V}_1 w\right| \le 2M\max|w|$. The boundedness of $\mathbf{V}_2$ is proved analogously.

The inverse operators $\mathbf{V}_1^{-1}$ and $\mathbf{V}_2^{-1}$ have the form

$$\mathbf{V}_1^{-1} = T_f^{-1}\mathcal{R} + \mathbf{j}T_{1/f}^{-1}\mathcal{I} \qquad \text{and} \qquad \mathbf{V}_2^{-1} = T_{1/f}^{-1}\mathcal{R} + \mathbf{j}T_f^{-1}\mathcal{I}$$

and clearly are bounded. For the explicit formulae for T_f^{-1} and $T_{1/f}^{-1}$ we refer to [**26**, Theorem 10].

The following statement establishes a relation between the derivatives of hyperbolic analytic functions and the generalized derivatives of their images under the action of the transmutation operator.

PROPOSITION 5.3. *Let w be a hyperbolic analytic function in $\mathbf{R}$ and $W = \mathbf{V}_1 w$ be a corresponding solution of* (2.4). *Then whenever the corresponding derivative of w exists there* *exists a derivative of the same order in the sense of Bers of the function W and vice versa, and the following relations are valid*

$$\mathbf{V}_1\left(\partial_z^{(2n)}w\right)=W^{[2n]}\quad \textit{and}\quad \mathbf{V}_2\left(\partial_z^{(2n-1)}w\right)=W^{[2n-1]},\quad n=1,2,\ldots. \tag{5.3}$$

Proof. From Proposition 5.1 we have

$$\overset{\circ}{W}=\mathbf{V}_2\left(\partial_z w\right). \tag{5.4}$$

$\overset{\circ}{W}$ is a solution of the succeeding Vekua equation (2.10). Denote $W_1=\overset{\circ}{W}$. Any solution of (2.10) is the image of a bicomplex analytic function under the action of the operator $\mathbf{V}_2$, so $W_1=\mathbf{V}_2w_1$. Due to Proposition 5.1 we have $\overset{\circ}{W}_1=\mathbf{V}_1\left(\partial_z w_1\right)$. Thus, $\overset{\circ\circ}{W}=\mathbf{V}_1\left(\partial_z^2 w\right)$ because from (5.4) $w_1=\partial_z w$. Now (5.3) can be easily proved by induction. □

6. *Expansion theorem*

Theorem 6.1. *A solution W of* (2.5) *can be represented as a uniformly convergent series*

$$W(z)=u(z)+\mathbf{j}v(z)=\sum_{n=0}^{\infty}Z^{(n)}(a_n,0;z) \tag{6.1}$$

in $\overline{\mathbf{R}}$ iff the functions $u(x,x)$ and $u(x,-x)$ admit the uniformly convergent series expansions on $[-b,b]$ of the form (4.3) *and* (4.4) *where the coefficients of the expansions are related by the equalities*

$$a_n=c_n+\mathbf{j}b_n,\quad n=0,1,\ldots$$

and the coefficient b_0 defines the value of v at the origin.

Proof. Suppose that W is a solution of (2.5) such that (6.1) holds. Denote $c_n:=\mathcal{R}(a_n)$ and $b_n:=\mathcal{I}(a_n)$, $n=0,1,\ldots$. By (2.28) we have

$$Z^{(n)}(1,0;z)=u_{2n-1}(x,t)+\mathbf{j}v_{2n}(x,t),\quad n=1,2,\ldots$$

and

$$Z^{(n)}(\mathbf{j},0;z)=u_{2n}(x,t)+\mathbf{j}v_{2n-1}(x,t),\quad n=1,2,\ldots.$$

Then

$$W(z)=c_0u_0(x,t)+b_0\mathbf{j}v_0(x,t)+\sum_{n=1}^{\infty}\left(c_n\left(u_{2n-1}(x,t)+\mathbf{j}v_{2n}(x,t)\right)+b_n\left(u_{2n}(x,t)+\mathbf{j}v_{2n-1}(x,t)\right)\right).$$

Hence

$$u:=\mathcal{R}(W)=c_0u_0+\sum_{n=1}^{\infty}\left(c_nu_{2n-1}+b_nu_{2n}\right)$$

and

$$v:=\mathcal{I}(W)=b_0v_0+\sum_{n=1}^{\infty}\left(c_nv_{2n}+b_nv_{2n-1}\right).$$

Thus,

$$u(x,x)=c_0u_0(x,x)+\sum_{n=1}^{\infty}\left(c_nu_{2n-1}(x,x)+b_nu_{2n}(x,x)\right)$$

and

$$u(x,-x)=c_0u_0(x,x)+\sum_{n=1}^{\infty}\left(c_nu_{2n-1}(x,x)-b_nu_{2n}(x,x)\right).$$

We obtain that (4.3) and (4.4) hold.

Now let us assume that W is a solution of (2.5) in $\overline{\mathbf{R}}$ such that for $u=\mathcal{R}(W)$ the equalities (4.3) and (4.4) hold. Then due to Proposition 4.1 the function u admits a series expansion (4.5), uniformly convergent in $\overline{\mathbf{R}}$ and at the same time $\widetilde{u}=T_f^{-1}u$ is a solution of a Goursat problem for the wave equation with the Goursat data $\binom{\varphi}{\psi}$ defined by (4.6). $\widetilde{u}$ has the form (4.7).

Now, let us consider the Goursat problem (2.33), (2.35) with

$$\Phi(x):=2\varphi(x)-c_0+b_0 \quad \text{and} \quad \Psi(x):=2\psi(x)-c_0-b_0$$

where $b_0\in\mathbb{C}$ is an arbitrary number. Thus,

$$\Phi(x)=\sum_{n=0}^{\infty}2^n\left(c_n+b_n\right)x^n \quad \text{and} \quad \Psi(x)=\sum_{n=0}^{\infty}2^n\left(c_n-b_n\right)x^n.$$

Due to Proposition 2.6, the unique solution of this Goursat problem has the form $\widetilde{W}(z)=\sum\limits_{n=0}^{\infty}a_nz^n$ where

$$a_n=P^+\left(c_n+b_n\right)+P^-\left(c_n-b_n\right)=c_n+\mathbf{j}b_n,\quad n=0,1,\ldots.$$

Application of the operator $\mathbf{V}_1$ to $\widetilde{W}$ gives us a solution W of (2.5) in $\overline{\mathbf{R}}$ in the form of a uniformly convergent series (6.1) with $u=\mathcal{R}(W)$. □

THEOREM 6.2. *Let W be a solution of* (2.5) *admitting in $\overline{\mathbf{R}}$ a uniformly convergent series expansion* (6.1). *Then there exist its derivatives in the sense of Bers of any order and the Taylor formula for the expansion coefficients holds,*

$$a_n=\frac{W^{[n]}(0)}{n!}. \tag{6.2}$$

Proof. Under the condition of the theorem we obtain that $w=\mathbf{V}_1^{-1}W$ has the form (2.36) with the expansion coefficients (2.38). The derivatives of w of any order exist because they reduce to the differentiation of power series expansions of Φ and Ψ from Proposition 2.6. Application of the operator $\mathbf{V}_1$ to the even derivatives and of $\mathbf{V}_2$ to the odd, due to Proposition 5.3 gives us the corresponding derivatives of W in the sense of Bers as well as the formula (6.2). □

Even if the conditions of Theorem 6.1 are not fulfilled and a solution of (2.5) cannot be represented globally as a uniformly convergent formal power series of the form (6.1), it may be arbitrarily closely approximated by finite linear combinations of the formal powers. The following analogue of the Runge approximation theorem from classical complex analysis is a corollary of the existence of the transmutation operator $\mathbf{V}_1$.

PROPOSITION 6.3. *Let W be a solution of* (2.5) *in* $\mathbf{R}$. *Then there exists a sequence of polynomials in formal powers $P_N=\sum_{n=0}^{N}Z^{(n)}(a_n,0;z)$ uniformly convergent to W in $\overline{\mathbf{R}}$.*

Proof. Consider $w=\mathbf{V}_1^{-1}W$. By Proposition 5.1 we have $\mathbf{V}_2(\partial_{\bar z}w)=0$, hence w is a solution of (2.33) in $\overline{\mathbf{R}}$. Despite the domain $\overline{\mathbf{R}}$ is smaller than the domain $\overline{R}$, the proof of

Proposition 2.10 may be applied without changes to the function w. Hence for an arbitrary $\varepsilon > 0$ there exists a polynomial $p_n = \sum_{n=0}^{N_\varepsilon} a_n z^n$ such that for any $(x,t) \in \overline{\mathbf{R}}$

$$|w(x,t) - p_n(x,t)| < \frac{\varepsilon}{\|\mathbf{V}_1\|},$$

where $\|\mathbf{V}_1\| < \infty$ is the norm of the transmutation operator $\mathbf{V}_1$. Applying $\mathbf{V}_1$ and Proposition 5.2 we obtain that

$$\left| W(x,t) - \sum_{n=0}^{N_\varepsilon} Z^{(n)}(a_n, 0; z) \right| \le \|\mathbf{V}_1\| \cdot \max_{(x,t)\in\overline{\mathbf{R}}} |w(x,t) - p_n(x,t)| < \varepsilon.$$

□

7. *Integral kernels as solutions of Vekua equations*

In Section 5 we showed that combinations (5.1) and (5.2) of the transmutation operators T_f and $T_{1/f}$ are related with Vekua operators and formal powers. In this section we consider the combination

$$\mathbf{K} = \mathbf{K}_f - \mathbf{j}\mathbf{K}_{1/f} \tag{7.1}$$

of the integral kernels of the operators T_f and $T_{1/f}$.

Theorem 7.1. *The function* $\mathbf{K}$ *given by* (7.1) *is a solution of the hyperbolic Vekua equation* (2.5).

Proof. The proof immediately follows from relations (3.11) and (3.12) and conditions (2.6), (2.7). □

The way of constructing the integral kernel $\mathbf{K}_{1/f}$ by the integral kernel $\mathbf{K}_f$ given by (3.11) requires the knowledge of the kernel $\mathbf{K}_f(x,t)$ not only on a natural domain $|t| \le |x| \le b$ sufficient for defining the transmutation operator, but on a larger domain $|t| \le b$, $|x| \le b$. Pseudoanalytic function theory allows us to obtain another method of reconstructing $\mathbf{K}_{1/f}$ by the known $\mathbf{K}_f$ and vice versa. Resulted formulae are simpler than those mentioned in [**26**, Remark 21].

Corollary 7.2. *The integral kernels* $\mathbf{K}_f$ *and* $\mathbf{K}_{1/f}$ *of the transmutation operators* T_f *and* $T_{1/f}$ *are related by the expressions*

$$\mathbf{K}_{1/f}(x,t) = -\frac{1}{f(x)}\int_0^x f(\eta)\partial_t \mathbf{K}_f(\eta,0)\,d\eta \\ - \int_0^t \partial_x \mathbf{K}_f(x,\xi)\,d\xi + \frac{f'(x)}{f(x)}\int_0^t \mathbf{K}_f(x,\xi)\,d\xi - \frac{f'(0)}{2f(x)} \tag{7.2}$$

and

$$\mathbf{K}_f(x,t) = -f(x)\int_0^x \frac{1}{f(\eta)}\partial_t \mathbf{K}_{1/f}(\eta,0)\,d\eta \\ - \int_0^t \partial_x \mathbf{K}_{1/f}(x,\xi)\,d\xi - \frac{f'(x)}{f(x)}\int_0^t \mathbf{K}_{1/f}(x,\xi)\,d\xi + \frac{f'(0)}{2}f(x). \tag{7.3}$$

Proof. Suppose first that $q_f := \frac{f''}{f} \in C^1[-b,b]$. In such case we also have $q_{1/f} = 2\Big(\frac{f'}{f}\Big)^2 - q_f \in C^1[-b,b]$ and the integral kernels $\mathbf{K}_f$ and $\mathbf{K}_{1/f}$ are twice continuously differentiable with respect to both variables and satisfy equation (3.5) with potentials q_f and $q_{1/f}$ correspondingly. Hence we obtain from Theorems 2.2 and 7.1 for the function $\mathbf{K}_f$ using the operator $\overline{A}_h$ given by (2.14) that the function $\mathbf{K}_{1/f}$ differs from the function

$$-v = -\frac{1}{f(x)}\int_0^x f(\eta)\partial_t \mathbf{K}_f(\eta,0)\,d\eta - \int_0^t \partial_x \mathbf{K}_f(x,\xi)\,d\xi + \frac{f'(x)}{f(x)}\int_0^t \mathbf{K}_f(x,\xi)\,d\xi$$

by the term cf^{-1}, where c is a constant. From the condition $\mathbf{K}_{1/f}(0,0) = -\frac{f'(0)}{2}$ (condition (3.6)) we find that $c = -\frac{f'(0)}{2}$.

Suppose now that $q_f \in C[-b,b]$. We proceed as follows. Consider a sequence of polynomials $\{f_n\}_{n\in\mathbb{N}}$ such that $f_n(0) = 1$ and $f_n'' \to f''$, $f_n' \to f'$ and $f_n \to f$ uniformly on $[-b,b]$ as $n\to\infty$. See, e.g., [**6**, Proof of Theorem 11] for a possible construction. Since the function f does not vanish on $[-b,b]$ we may additionally assume that all functions f_n do no vanish on $[-b,b]$. It is easy to see that $q_n := \frac{f_n''}{f_n} \to q$ uniformly and $q_n \in C^1[-b,b]$. Every f_n is associated with the transmutation operators T_{f_n} and T_{1/f_n} with the integral kernels $\mathbf{K}_{f_n}$ and $\mathbf{K}_{1/f_n}$. Consider functions $\mathbf{H}_{f_n}(u,v) = \mathbf{K}_{f_n}(u+v, u-v; f_n'(0))$. It is well known (e.g., [**46**]) that the Goursat problem (3.3), (3.4) is equivalent to either of the systems of integral equations

$$\begin{cases}\mathbf{H}(u,v) = \frac{h}{2} + \int_0^u \mathbf{G}(u',v)\,du',\\ \mathbf{G}(u,v) = \frac{1}{2}q(u) + \int_0^v q(u+v')\mathbf{H}(u,v')\,dv'\end{cases}$$

and

$$\begin{cases}\mathbf{H}(u,v) = \frac{h}{2} + \frac{1}{2}\int_0^u q(s)\,ds + \int_0^v \mathbf{F}(u,v')\,dv',\\ \mathbf{F}(u,v) = \int_0^u q(u'+v)\mathbf{H}(u',v)\,du',\end{cases}$$

where $\mathbf{G} = \partial_u \mathbf{H}$ and $\mathbf{F} = \partial_v\mathbf{H}$, from which we conclude that the functions $\mathbf{H}_{f_n}$, $\partial_u\mathbf{H}_{f_n}$ and $\partial_v\mathbf{H}_{f_n}$ are uniformly convergent to the functions $\mathbf{H}_f$, $\partial_u\mathbf{H}_f$ and $\partial_v\mathbf{H}_f$, respectively. Hence the partial derivatives of $\mathbf{K}_{f_n}$ are uniformly convergent as well, and taking in (7.2) the limit as $n\to\infty$ we finish the proof.

The second formula (7.3) easily follows from (7.2) by changing f to $1/f$. □

Example 7.3. For the functions $\mathbf{K}_f$ and $\mathbf{K}_{1/f}$ from Example 3.10 we get $\mathbf{K} = \frac{1}{2} - \mathbf{j}\frac{t-1}{2(x+1)}$ and

$$\partial_{\bar z}\mathbf{K} = \frac{1}{2}\left(\mathbf{j}\frac{t-1}{2(x+1)^2} + \frac{1}{2(x+1)}\right) = \frac{1}{2(x+1)}\overline{\mathbf{K}} = \frac{f'(x)}{2f(x)}\overline{\mathbf{K}}.$$

Formulae (7.2) and (7.3) can be verified as well.

8. *Formulae for integral kernels of transmutation operators*

It follows from Theorems 6.2 and 7.1 and from (2.12) that the knowledge of the derivatives $\partial_t^n\mathbf{K}(0,0)$ allows one to obtain the generalized Taylor coefficients (6.2) for $W = \mathbf{K}$. In this section we present several formulae for computing these derivatives whenever the derivatives $q^{(m)}(0)$, $m \le n-1$ are known.

First we illustrate the expansion theorem with an example.

Example 8.1. For the function $\mathbf{K}$ from Example 7.3 by (2.12) we obtain

$$\mathbf{K}^{[0]}(0,0)=\frac{1}{2}+\frac{1}{2}\mathbf{j},\quad \mathbf{K}^{[1]}(0,0)=-\frac{1}{2(x+1)}\Big|_{(0,0)}=-\frac{1}{2}+0\mathbf{j},\quad \mathbf{K}^{[n+1]}=\mathbf{j}\partial_t\mathbf{K}^{[n]}=0,\ n\geq 1.$$

Hence the expansion coefficients are $a_0=\frac{1}{2}+\frac{1}{2}\mathbf{j}$, $a_1=-\frac{1}{2}+0\mathbf{j}$ and $a_n=0$ for $n\geq 2$.

The first two functions of the systems $\{\varphi_k\}$ and $\{\psi_k\}$ are equal to

$$\varphi_0=x+1,\qquad \psi_0=\frac{1}{x+1},$$
$$\varphi_1=x,\qquad \psi_1=\frac{x^3+3x^2+3x}{3(x+1)},$$

the first two formal powers are given by

$$Z^{(0)}(a,0;z)=\mathcal{R}(a)\cdot\varphi_0+\mathbf{j}\mathcal{I}(a)\cdot\psi_0,$$
$$Z^{(1)}(a,0;z)=\mathcal{R}(a)\cdot(\varphi_1+\mathbf{j}t\psi_0)+\mathbf{j}\mathcal{I}(a)\cdot(\psi_1+\mathbf{j}t\varphi_0)$$

and we find that indeed,

$$\mathbf{K}(x,t)=Z^{(0)}\Big(\frac{1}{2}+\frac{1}{2}\mathbf{j},0;z\Big)+Z^{(1)}\Big(-\frac{1}{2}+0\mathbf{j},0;z\Big).$$

We are able to calculate the expansion coefficients a_n whenever we know values of the derivatives $\partial_t^n\mathbf{K}(0,0)$. Recall that the function $\mathbf{K}_f$ is the integral kernel of a transmutation operator if and only if the function $\mathbf{H}_f(u,v)=\mathbf{K}_f(u+v,u-v)$ is the solution of the Goursat problem (3.3), (3.4). We have

$$\partial_t^n\mathbf{K}_f(x,t)=\partial_t^n\mathbf{H}_f\Big(\frac{x+t}{2},\frac{x-t}{2}\Big)=\frac{1}{2^n}(\partial_u-\partial_v)^n\mathbf{H}_f(u,v)\Big|_{u=\frac{x+t}{2},\,v=\frac{x-t}{2}}.\tag{8.1}$$

Moreover, it follows from (3.4) that

$$\partial_u^n\mathbf{H}_f(u,0)=\begin{cases}\frac{f'(0)}{2}+\frac{1}{2}\int_0^u q_f(s)\,ds, & n=0,\\ \frac{1}{2}q_f^{(n-1)}(u), & n\geq 1,\end{cases}\tag{8.2}$$

and

$$\partial_v^n\mathbf{H}_f(0,v)=\begin{cases}\frac{f'(0)}{2}, & n=0,\\ 0, & n\geq 1.\end{cases}\tag{8.3}$$

Hence to calculate the value of $\partial_t^n\mathbf{K}_f(0,0)$ it is sufficient to transform the terms $\partial_u^k\partial_v^{n-k}\mathbf{H}_f$, where $1\leq k\leq n-1$, into the terms involving only derivatives of the form $\partial_u^\ell\mathbf{H}_f$ or $\partial_v^m\mathbf{H}_f$ and some derivatives of the potential q_f. Such transformation may be performed using equation (3.3) which allows us to reduce the orders k and $n-k$ in the term $\partial_u^k\partial_v^{n-k}\mathbf{H}_f$ by 1. After several applications of equation (3.3) and the Leibniz rule we obtain a finite sum of terms of the form

$$q_f^{(n_1)}(u+v)\cdot\ldots\cdot q_f^{(n_\ell)}(u+v)\partial_{u/v}^d\mathbf{H}_f(u,v).\tag{8.4}$$

It is easy to see that the number ℓ of factors $q^{(n_i)}(u+v)$ satisfies the inequality $1\leq\ell\leq\min(k,n-k)$ and that $n_1+\ldots+n_\ell+2\ell+d=n$.

Moreover, consider a term of the form (8.4) obtained as a result of application to $\mathbf{H}_f$ of some derivatives ∂_u and ∂_v, equation (3.3) and the Leibniz rule. If during this process we change every derivative ∂_u by $-\partial_v$ and every derivative ∂_v by $-\partial_u$, we obtain the term of exactly the same form, with the only difference that the last derivative $\partial_{u/v}^d\mathbf{H}_f$ changes into $\partial_{v/u}^d\mathbf{H}_f$, and the coefficient $(-1)^n$ appears. Hence, we may group all the terms obtained after the transformation of the expression $(\partial_u-\partial_v)^n\mathbf{H}_f(u,v)$ into the pairs of the form

$$\big(\partial_u^d+(-1)^n\partial_v^d\big)\mathbf{H}_f(u,v)\cdot\prod_{i=1}^{\ell}q_f^{(n_i)}(u+v). \tag{8.5}$$

Now we derive the recurrent relation for the derivative $(\partial_u-\partial_v)^n\mathbf{H}_f$. We may parametrize each term of the form (8.5) by the following parameters.

(1) A number ℓ of the factors $q_f^{(n_i)}(u+v)$. Since we need two derivatives to obtain one factor $q(u+v)$ with the use of (3.3), the number ℓ satisfies

$$0\le\ell\le\frac{n}{2}. \tag{8.6}$$

(2) A number d of derivatives of $\mathbf{H}_f$ satisfying

$$0\le d\le n-2\ell. \tag{8.7}$$

(3) An ordered sequence of non-negative numbers $n_1\le n_2\le\ldots\le n_\ell$ (orders of derivatives of the factors $q_f(u+v)$) satisfying

$$n_1+n_2+\ldots+n_\ell+2\ell+d=n. \tag{8.8}$$

We denote by $(-1)^\ell S^n_{\ell;d;(n_1,\ldots,n_\ell)}$ the coefficient at the term (8.5) in the final representation of $(\partial_u-\partial_v)^n\mathbf{H}_f$ after the described transformation. The factor $(-1)^\ell$ is used to make the number $S^n_{\ell;d;(n_1,\ldots,n_\ell)}$ non-negative. We assume that $S^n_{\ell;d;(n_1,\ldots,n_\ell)}=0$ whenever either of the conditions (8.6)–(8.8) fails.

Lemma 8.2. *The number of different lists of parameters $\ell;d;(n_1,\ldots,n_\ell)$ satisfying conditions* (8.6)–(8.8) *for a fixed n coincides with the number $p(n)$ of partitions of n, i.e., the number of ways n may be represented as a sum of ordered positive integer terms, see, e.g.,* [**10**], [**11**]. *The number $p(n)$ is also known as the number of different Young or Ferrers diagrams.*

Proof. Consider a list of parameters $\ell;d;(n_1,\ldots,n_\ell)$ and put into correspondence to this list a partition $\underbrace{1,1,\ldots,1}_{d},n_1+2,n_2+2,\ldots,n_\ell+2$. Due to the condition (8.8) this partition is a partition of n. It is easy to see that the described correspondence is one-to-one. □

Note that $(\partial_u-\partial_v)q_f(u+v)=0$, hence in the case $d=0$ we have

$$\begin{aligned}(\partial_u-\partial_v)\Big(\big(\mathbf{H}_f(u,v)+(-1)^n\mathbf{H}_f(u,v)\big)\prod_{i=1}^{\ell}q_f^{(n_i)}(u+v)\Big)&\\ =\big(1+(-1)^n\big)\big(\partial_u+(-1)^{n+1}\partial_v\big)\mathbf{H}_f(u,v)\cdot\prod_{i=1}^{\ell}q_f^{(n_i)}(u+v),&\end{aligned} \tag{8.9}$$

and in the case $d\ne0$ we have

$$\begin{aligned}&(\partial_u-\partial_v)\Big(\big(\partial_u^d+(-1)^n\partial_v^d\big)\mathbf{H}_f(u,v)\prod_{i=1}^{\ell}q_f^{(n_i)}(u+v)\Big)\\ &=\Big(\big(\partial_u^{d+1}+(-1)^{n+1}\partial_v^{d+1}\big)\mathbf{H}_f(u,v)-\big(\partial_u^{d-1}+(-1)^{n+1}\partial_v^{d-1}\big)\big(q_f(u+v)\mathbf{H}_f(u,v)\big)\Big)\prod_{i=1}^{\ell}q_f^{(n_i)}(u+v)\\ &=\big(\partial_u^{d+1}+(-1)^{n+1}\partial_v^{d+1}\big)\mathbf{H}_f(u,v)\cdot\prod_{i=1}^{\ell}q_f^{(n_i)}(u+v)\end{aligned}$$

$$-\sum_{k=0}^{d-1}\binom{d-1}{k}q_f^{(k)}(u+v)\big(\partial_u^{d-1-k}+(-1)^{n+1}\partial_v^{d-1-k}\big)\mathbf{H}_f(u,v)\cdot\prod_{i=1}^{\ell}q_f^{(n_i)}(u+v). \tag{8.10}$$

Also note that the coefficient at the term $(\partial_u^n+(-1)^n\partial_v^n)\mathbf{H}_f(u,v)$ is equal to $1/2$ for $n=0$ and 1 for $n\geq 1$, hence $S^0_{0;0;()}=1/2$ and $S^n_{0;n;()}=1$ for $n\geq 1$. From the relations (8.9) and (8.10) and the latter initial condition we obtain the following statement.

Proposition 8.3. *The coefficients* $S^{n+1}_{\ell;d;(n_1,\dots,n_\ell)}$ *with the lists of parameters* $\ell;d;(n_1,\dots,n_\ell)$ *satisfying conditions* (8.6)–(8.8) *can be calculated by means of the following recurrent relations*

$$\begin{aligned}S^{n+1}_{\ell;d;(n_1,\dots,n_\ell)}=&\big(1+(-1)^n\big)S^n_{\ell;d-1;(n_1,\dots,n_\ell)}\\&+\sum_{n_k\in\{n_1,\dots,n_\ell\}}\binom{d+n_k}{n_k}S^n_{\ell-1;d+n_k+1;(n_1,\dots,n_{k-1},n_{k+1},\dots,n_\ell)}\qquad\text{if } d=1,\end{aligned}\tag{8.11}$$

$$\begin{aligned}S^{n+1}_{\ell;d;(n_1,\dots,n_\ell)}=&S^n_{\ell;d-1;(n_1,\dots,n_\ell)}\\&+\sum_{n_k\in\{n_1,\dots,n_\ell\}}\binom{d+n_k}{n_k}S^n_{\ell-1;d+n_k+1;(n_1,\dots,n_{k-1},n_{k+1},\dots,n_\ell)}\qquad\text{if } d\neq 1,\end{aligned}\tag{8.12}$$

where $\sum_{n_k\in\{n_1,\dots,n_\ell\}}$ *means that only different values of* n_i *are used in the sum, with the initial conditions*

$$S^0_{0;0;()}=1/2,\qquad S^n_{0;n;()}=1\quad\text{for } n\geq 1. \tag{8.13}$$

The following proposition presents a direct formula for the coefficients $S^n_{\ell;d;(n_1,\dots,n_\ell)}$ not involving recurrent relations.

Proposition 8.4. *The following formula holds for the coefficient* $S^n_{\ell;d;(n_1,\dots,n_\ell)}$, *where* $\ell\geq 1$ *and the list of parameters* $\ell;d;(n_1,\dots,n_\ell)$ *satisfies conditions* (8.6)–(8.8)*:*

$$\begin{aligned}S^n_{\ell;d;(n_1,\dots,n_\ell)}=&\sum_{(\sigma_1,\dots,\sigma_\ell)\in\mathfrak{S}(n_1,\dots,n_\ell)}\sum_{d_1=0}^{d}\sum_{d_2=0}^{d_1+\sigma_1+1}\cdots\sum_{d_\ell=0}^{d_{\ell-1}+\sigma_{\ell-1}+1}\\&\prod_{i=1}^{\ell}\binom{d_i+\sigma_i}{\sigma_i}\big(1+(-1)^{\sigma_i+\ldots+\sigma_\ell}\delta(d_i)\big(1-\delta(d)\delta(i-1)\big)\big),\end{aligned}\tag{8.14}$$

where $(\sigma_1,\dots,\sigma_\ell)\in\mathfrak{S}(n_1,\dots,n_\ell)$ *means that the sum is taken over all distinct permutations of the numbers* $n_1,\dots,n_\ell$, *and* $\delta(t)=1$ *if* $t=0$ *and* $\delta(t)=0$ *otherwise.*

Proof. To arrive from the term $S^n_{\ell;d;(n_1,\dots,n_\ell)}$ to a term $S^{\tilde n}_{0;\tilde n;()}$ using recurrent relations (8.11) and (8.12) we have to perform exactly ℓ times the step of removing one of the n_k's from the list $(n_1,\dots,n_\ell)$. Denote by $d_1,\dots,d_\ell$ the values of the parameter d on each of these steps and by $\sigma_1,\dots,\sigma_\ell$ the number n_k removed from the list $(n_1,\dots,n_\ell)$ on the corresponding step $1,\dots,\ell$. Then the numbers $d_1,\dots,d_\ell$ satisfy the following inequalities

$$\begin{aligned}&0\leq d_1\leq d,\\&0\leq d_2\leq 1+d_1+\sigma_1,\\&\qquad\vdots\\&0\leq d_\ell\leq 1+d_{\ell-1}+\sigma_{\ell-1}.\end{aligned}$$

Each distinct permutation of the numbers $n_1,\ldots,n_\ell$ and each list of parameters $d_1,\ldots,d_\ell$ give us a different way of getting from the term $S^n_{\ell;d;(n_1,\ldots,n_\ell)}$ to the term $S^{\tilde n}_{0;\tilde n;()}$. Note that the factor $1+(-1)^{n'}$ for some $n'\le n$ appears in (8.11) only in the case when we cross the coefficient $S^{n'}_{\ell';0;(n_1',\ldots,n_{\ell'}')}$ having $d'=0$. Note also that if $d=0$ initially, then $d_1=0$ and only (8.12) is applicable, hence the factor $1+(-1)^n$ does not appear in this case. So the factor $1+(-1)^{n'}$ appears only in one of the following cases: 1) $d\ne 0$ and $d_1=0$, 2) $d_i=0$, $i\ge 2$. It follows from condition (8.8) that in each of these cases $n'=2\ell'+n_1'+\ldots+n_{\ell'}'$ therefore $(-1)^{n'}=(-1)^{n_1'+\cdots+n_{\ell'}'}$. Hence we conclude that the following relation holds

$$S^n_{\ell;d;(n_1,\ldots,n_\ell)}=\sum_{(\sigma_1,\ldots,\sigma_\ell)\in\mathfrak{S}(n_1,\ldots,n_\ell)}\sum_{d_1=0}^{d}\binom{d_1+\sigma_1}{\sigma_1}\bigl(1+(-1)^{\sigma_1+\ldots+\sigma_\ell}\cdot\delta(d_1)(1-\delta(d))\bigr)$$
$$\times\sum_{d_2=0}^{d_1+\sigma_1+1}\binom{d_2+\sigma_2}{\sigma_2}\bigl(1+(-1)^{\sigma_2+\ldots+\sigma_\ell}\delta(d_2)\bigr)\cdots\sum_{d_\ell=0}^{d_{\ell-1}+\sigma_{\ell-1}+1}\binom{d_\ell+\sigma_\ell}{\sigma_\ell}\bigl(1+(-1)^{\sigma_\ell}\delta(d_\ell)\bigr),$$

where the first sum is taken over all distinct permutations of numbers $n_1,\ldots,n_\ell$, and the function $\delta(\cdot)$ is used to the encode mentioned above conditions 1) and 2). Finally note that the conditions 1) and 2) may be jointly represented by the expression $\delta(d_i)\bigl(1-\delta(d)\delta(i-1)\bigr)$. $\square$

As a result we obtain the following representation for the derivative $(\partial_u-\partial_v)^n\mathbf{H}_f(u,v)$, $n\ge 1$.

Proposition 8.5. *Let $n\ge 1$ and $q_f\in C^{(n-1)}[-b,b]$. Then*

$$(\partial_u-\partial_v)^n\mathbf{H}_f(u,v)=\bigl(\partial_u^n+(-1)^n\partial_v^n\bigr)\mathbf{H}_f(u,v)$$
$$+\sum_{\ell=1}^{[n/2]}(-1)^\ell\sum_{d=0}^{n-2\ell}\bigl(\partial_u^d+(-1)^n\partial_v^d\bigr)\mathbf{H}_f(u,v)\sum_{\substack{n_1+\ldots+n_\ell=n-2\ell-d,\\ 0\le n_1\le\ldots\le n_\ell}}S^n_{\ell;d;(n_1,\ldots,n_\ell)}\prod_{i=1}^{\ell}q_f^{(n_i)}(u+v),$$

where the coefficients $S^n_{\ell;d;(n_1,\ldots,n_\ell)}$ are given by Proposition 8.3 or Proposition 8.4 and do not depend on q_f.

The following proposition is a corollary of (8.1), (8.2), (8.3) and Proposition 8.5.

Proposition 8.6. *Let $n\ge 1$ and $q_f\in C^{(n-1)}[-b,b]$. Then*

$$\partial_t^n\mathbf{K}_f(0,0)=\frac{1}{2^{n+1}}\Biggl(q_f^{(n-1)}(0)+\sum_{\ell=1}^{[n/2]}(-1)^\ell\sum_{d=0}^{n-2\ell}\bigl(1+(-1)^n\delta(d)\bigr)q_f^{(d-1)}(0)$$
$$\times\sum_{\substack{n_1+\ldots+n_\ell=n-d-2\ell,\\ 0\le n_1\le\ldots\le n_\ell}}S^n_{\ell;d;(n_1,\ldots,n_\ell)}\prod_{i=1}^{\ell}q_f^{(n_i)}(0)\Biggr),\qquad(8.15)$$

where we set $q_f^{(-1)}(0):=h=f'(0)$, the coefficients $S^n_{\ell;d;(n_1,\ldots,n_\ell)}$ are given by Proposition 8.3 or Proposition 8.4 and do not depend on q_f, and $\delta(d)=1$ if $d=0$ and $\delta(d)=0$ otherwise.

Formula (8.15) also holds for $n=0$ for any $q_f\in C[-b,b]$.

As a consequence of Theorems 6.1, 6.2 and 7.1 we obtain the following representation of the integral kernel $\mathbf{K}_f$.

THEOREM 8.7. *Let $q_f \in C^\infty[-b,b]$ be a complex-valued function and f be a particular solution of (2.30) such that $f \neq 0$ on $[-b,b]$ and normalized as $f(0)=1$. Denote $h := f'(0) \in \mathbb{C}$. Suppose that the functions $g_1(x) := \frac{1}{2}\big(\mathbf{K}_f(x,x)+\mathbf{K}_f(x,-x)\big) = \frac{h}{2}+\frac{1}{4}\int_0^x q_f(s)\,ds$ and $g_2(x) := \frac{1}{2}\big(\mathbf{K}_f(x,x)-\mathbf{K}_f(x,-x)\big) = \frac{1}{4}\int_0^x q_f(s)\,ds$ admit uniformly convergent on $[-b,b]$ series expansions*

$$g_1(x) = c_0u_0(x,x) + \sum_{n=1}^{\infty} c_n u_{2n-1}(x,x)$$

and

$$g_2(x) = \sum_{n=1}^{\infty} b_n u_{2n}(x,x).$$

Then the coefficients $\{c_n\}_{n\ge 0}$ and $\{b_n\}_{n\ge 1}$ may be found by the formulas

$$c_n = \frac{\partial_t^n \mathbf{K}_f(0,0)}{n!}, \qquad b_n = -\frac{\partial_t^n \mathbf{K}_{1/f}(0,0)}{n!}, \quad \text{if } n \text{ is even}$$

and

$$c_n = -\frac{\partial_t^n \mathbf{K}_{1/f}(0,0)}{n!}, \qquad b_n = \frac{\partial_t^n \mathbf{K}_f(0,0)}{n!}, \quad \text{if } n \text{ is odd,}$$

where the derivatives $\partial_t^n\mathbf{K}_f(0,0)$ and $\partial_t^n\mathbf{K}_{1/f}(0,0)$ are given by (8.15) for the potentials q_f and $q_{1/f} = -q_f + 2\big(f'/f\big)^2$, respectively. Also, for any $(x,t) \in \overline{\mathbf{R}}$

$$\mathbf{K}_f(x,t) = c_0u_0(x,t) + \sum_{n=1}^{\infty}\left(c_nu_{2n-1}(x,t) + b_nu_{2n}(x,t)\right), \tag{8.16}$$

and the series converges uniformly in $\overline{\mathbf{R}}$.

EXAMPLE 8.8. For the potentials q_f and $q_{1/f}$ from Example 3.10 we have

$$q_f^{(-1)}(0) := f'(0) = 1, \qquad q_f^{(n)}(0) = 0, \quad n \ge 0,$$

and

$$q_{1/f}^{(-1)}(0) := (1/f)'(0) = -1, \qquad q_{1/f}^{(n)}(0) = 2(-1)^n(n+1)!, \quad n \ge 0.$$

By Proposition 8.3 the first coefficients $S^n_{\ell;d;(n_1,\ldots,n_\ell)}$ for $n \le 6$ have the following values

$$S^1_{0;1;()} = 1;$$
$$S^2_{0;2;()} = 1, \quad S^2_{1;0;(0)} = 1;$$
$$S^3_{0;3;()} = 1; \quad S^3_{1;0;(1)} = 1, \quad S^3_{1;1;(0)} = 3;$$
$$S^4_{0;4;()} = 1; \quad S^4_{1;0;(2)} = 1, \quad S^4_{1;1;(1)} = 2; \quad S^4_{1;2;(0)} = 4, \quad S^4_{2;0;(0,0)} = 3;$$
$$S^5_{0;5;()} = 1; \ S^5_{1;0;(3)} = 1, \ S^5_{1;1;(2)} = 5; \ S^5_{1;2;(1)} = 5, \ S^5_{1;3;(0)} = 5, \ S^5_{2;0;(0,1)} = 6, \ S^5_{2;1;(0,0)} = 10;$$
$$S^6_{0;6;()} = 1; \quad S^6_{1;0;(4)} = 1, \quad S^6_{1;1;(3)} = 4; \quad S^6_{1;2;(2)} = 11, \quad S^6_{1;3;(1)} = 9, \quad S^6_{1;4;(0)} = 6,$$
$$S^6_{2;0;(0,2)} = 10, \quad S^6_{2;0;(1,1)} = 5, \quad S^6_{2;1;(0,1)} = 15, \quad S^6_{2;2;(0,0)} = 15, \quad S^6_{3;0;(0,0,0)} = 10.$$

We can check that formula (8.15) works, e.g.,

$$\partial_t^5\mathbf{K}_{1/f}(0,0) = \frac{1}{64}\Big(240 + (-1)^1\big(1\cdot(1+(-1)^5)\cdot 48 + 5\cdot 24 + 5\cdot 16 + 5\cdot 24\big) \\ + (-1)^2\big(6\cdot(1+(-1)^5)\cdot 8 + 10\cdot 8\big)\Big) = 0,$$

and

$$\begin{aligned}\partial_t^6\mathbf{K}_{1/f}(0,0) = \frac{1}{128}\Big(&-1440 + (-1)^1\big(1\cdot(1+(-1)^6)\cdot(-240) + 4\cdot(-96) + 11\cdot(-48)\\ &+ 9\cdot(-48) + 6\cdot(-96)\big) + (-1)^2\big(10\cdot(1+(-1)^6)\cdot(-24) + 5\cdot(1+(-1)^6)\cdot(-16)\\ &+ 15\cdot(-16) + 15\cdot(-16)\big) + (-1)^3\big(10\cdot(1+(-1)^6)\cdot(-8)\big)\Big) = 0.\end{aligned}$$

Now consider an example when all the coefficients b_n, c_n from (8.16) may be easily calculated explicitly from (8.14) and (8.15).

Example 8.9. Consider $q_f = c^2$ for some constant c taking the function $f = e^{cx}$ as a non-vanishing solution of the equation $f'' - q_f f = 0$. Then the Darboux transformed potential has the form $q_{1/f} = 2(f'/f)^2 - q_f = c^2$.

Consider the function $\mathbf{H}_f$ from (8.1). Equation (3.3) for $q_f = c^2$ reads as $\partial_u\partial_v\mathbf{H}_f = c^2\mathbf{H}_f$, hence

$$\begin{aligned}(\partial_u - \partial_v)^n\mathbf{H}_f(u,v) &= \sum_{k=0}^{n}(-1)^{n-k}\binom{n}{k}\partial_u^k\partial_v^{n-k}\mathbf{H}_f(u,v)\\ &= (\partial_u^n + (-1)^n\partial_v^n)\mathbf{H}_f(u,v)\\ &+ \sum_{k=1}^{[n/2]-1}(-1)^{n-k}c^{2k}\binom{n}{k}\partial_v^{n-2k}\mathbf{H}_f(u,v)\\ &+ \sum_{k=[n/2]}^{n-1}(-1)^{n-k}c^{2(n-k)}\binom{n}{k}\partial_u^{2k-n}\mathbf{H}_f(u,v).\end{aligned} \tag{8.17}$$

It follows from (8.2) and (8.3) that at the point $u = v = 0$ every term in (8.17) involving a derivative of $\mathbf{H}_f$ of order at least 2 with respect to u or at least 1 with respect to v equals zero. Hence from (8.1) for any even $n = 2m$ we obtain

$$\partial_t^{2m}\mathbf{K}_f(0,0) = \frac{1}{2^{2m}}(-1)^m c^{2m}\binom{2m}{m}\mathbf{H}_f(0,0) = \frac{(-1)^m c^{2m+1}}{2^{2m+1}}\binom{2m}{m}, \tag{8.18}$$

and for any odd $n = 2m+1$ we obtain

$$\partial_t^{2m+1}\mathbf{K}_f(0,0) = \frac{1}{2^{2m+1}}(-1)^m c^{2m}\binom{2m+1}{m}\partial_u\mathbf{H}_f(0,0) = \frac{(-1)^m c^{2m+2}}{2^{2m+2}}\binom{2m+1}{m}. \tag{8.19}$$

Similarly for the kernel $\mathbf{K}_{1/f}$ we have

$$\partial_t^{2m}\mathbf{K}_{1/f}(0,0) = -\frac{(-1)^m c^{2m+1}}{2^{2m+1}}\binom{2m}{m}, \qquad \partial_t^{2m+1}\mathbf{K}_{1/f}(0,0) = \frac{(-1)^m c^{2m+2}}{2^{2m+2}}\binom{2m+1}{m}.$$

Hence by Theorem 8.7 the expansion coefficients have the form

$$\begin{aligned} c_{2m} &= \frac{(-1)^m c^{2m+1}}{2^{2m+1}(m!)^2}, & b_{2m} &= \frac{(-1)^m c^{2m+1}}{2^{2m+1}(m!)^2},\\ c_{2m+1} &= \frac{(-1)^{m+1} c^{2m+2}}{2^{2m+2}m!(m+1)!}, & b_{2m+1} &= \frac{(-1)^m c^{2m+2}}{2^{2m+2}m!(m+1)!}.\end{aligned}$$

Now we show that it is possible to obtain the same coefficients from (8.14) and (8.15). Note that any term in (8.15) involving a positive order derivative of q_f is equal to zero. Hence $n_1 = n_2 = \ldots = n_\ell = 0$ and $n = 2\ell + d$. Moreover, $d \le 1$ and is equal either to 0 or 1 depending on the parity of n. So we have to calculate only one coefficient, $S^{2m}_{m;0;(0,\ldots,0)}$ for $n = 2m$ and

$S^{2m+1}_{m;1;(0,\dots,0)}$ for $n = 2m+1$. Formula (8.14) for the coefficient $S^{2m}_{m;0;(0,\dots,0)}$ takes the form

$$S^{2m}_{m;0;(0,\dots,0)} = \sum_{d_2=0}^{1}\big(1+\delta(d_2)\big)\sum_{d_3=0}^{d_2+1}\big(1+\delta(d_3)\big)\cdots\sum_{d_m=0}^{d_{m-1}+1}\big(1+\delta(d_m)\big).$$

We claim that

$$\sum_{d_{m-k+1}=0}^{d_{m-k}+1}\big(1+\delta(d_{m-k+1})\big)\cdots\sum_{d_m=0}^{d_{m-1}+1}\big(1+\delta(d_m)\big) = \binom{2k+1+d_{m-k}}{k}. \tag{8.20}$$

Indeed, for $k=1$ we have $\sum_{d_m=0}^{d_{m-1}+1}\big(1+\delta(d_m)\big) = d_{m-1}+3 = \binom{3+d_{m-1}}{1}$. Assume that the claim is correct for some k, then for $k+1$ we obtain

$$\sum_{d_{m-k}=0}^{d_{m-k-1}+1}\big(1+\delta(d_{m-k})\big)\cdot\binom{2k+1+d_{m-k}}{k}$$
$$= 2\binom{2k+1}{k}+\binom{2k+2}{k}+\ldots+\binom{2k+2+d_{m-k-1}}{k},$$

and the induction step follows from the equality $2\binom{2k+1}{k}+\binom{2k+2}{k}=\binom{2k+3}{k+1}$ and the well-known combinatorial identity $\binom{n+1}{k+1}=\binom{n}{k}+\binom{n-1}{k}+\ldots+\binom{n-s}{k}+\binom{n-s}{k+1}$, see, e.g., [**11**]. For $k=m-1$ we obtain from (8.20) that $S^{2m}_{m;0;(0,\dots,0)}=\binom{2m-1}{m-1}$ and $\partial_t^{2m}\mathbf{K}_f(0,0)=\frac{(-1)^m}{2^{2m+1}}2c\binom{2m-1}{m-1}(c^2)^m=\frac{(-1)^m c^{2m+1}}{2^{2m+1}}\binom{2m}{m}$.

Similarly, for $n=2m+1$ we obtain from (8.20) for $k=m$ that $S^{2m+1}_{m;1;(0,\dots,0)}=\binom{2m+1}{m}$ and $\partial_t^{2m+1}\mathbf{K}_f(0,0)=\frac{(-1)^m}{2^{2m+2}}c^2\binom{2m+1}{m+1}(c^2)^m=\frac{(-1)^m c^{2m+2}}{2^{2m+2}}\binom{2m+1}{m}$, exactly as in (8.18) and (8.19).

In the following example we calculate the expansion coefficients b_n and c_n, $n\le 22$, for a reflectionless potential in the one-dimensional quantum scattering theory, $q_{\text{sech}}(x)=1-2\operatorname{sech}^2 x$, see, e.g., [**26**], [**30**]. For this potential to the difference with from the previous examples neither the higher derivatives of q_f nor the higher derivatives with respect to t of the transmutation kernel do not vanish identically.

Example 8.10. The potential $q_{\text{sech}}(x)=1-2\operatorname{sech}^2 x$ can be obtained as a result of the Darboux transformation of the equation $u''=u$ with the potential $q_{\cosh}\equiv 1$ with respect to the solution $f(x)=\cosh x$. The transmutation operator for the operator $A_{\cosh}=\partial_x^2-1$ was calculated in [**6**, Example 3]. Its kernel is given by the expression

$$\mathbf{K}_{\cosh}(x,t)=\frac{1}{2}\frac{\sqrt{x^2-t^2}\,I_1(\sqrt{x^2-t^2})}{x-t},$$

where I_1 is the modified Bessel function of the first kind. Even though in [**26**] a formula for the integral kernel $\mathbf{K}_{\text{sech}}$ was presented, it is not well suited for calculating higher order derivatives with respect to t. Using (7.2) we obtain another representation with the help of Maple 12 software

$$\mathbf{K}_{\text{sech}}(x,t)=\frac{1}{2}\bigg(I_1(x)-I_0(x)\tanh x+\tanh x\int_0^t\frac{\sqrt{x^2-s^2}\,I_1(\sqrt{x^2-s^2})}{x-s}\,ds$$
$$+\int_0^t\frac{(xs-x^2)I_0(\sqrt{x^2-s^2})+\sqrt{x^2-s^2}\,I_1(\sqrt{x^2-s^2})}{(x-s)^2}\,ds\bigg). \tag{8.21}$$

Despite the integral kernel $\mathbf{K}_{\text{sech}}$ could not be evaluated explicitly, it is possible to expand it into a Taylor series at the point $t=0$. Expanding the kernels $\mathbf{K}_{\cosh}$ and $\mathbf{K}_{\text{sech}}$ into corresponding

Taylor series and taking a limit as $x \to 0$ we obtain (again with the help of Maple 12 software) that

$$\begin{aligned}\mathbf{K}_{\cosh}(0,t) = {}&\frac{1}{4}\,t - \frac{1}{32}\,t^3 + \frac{1}{768}\,t^5 - \frac{1}{36864}\,t^7 + \frac{1}{2949120}\,t^9 - \frac{1}{353894400}\,t^{11}\\ &+ \frac{1}{59454259200}\,t^{13} - \frac{1}{13317754060800}\,t^{15} + \frac{1}{3835513169510400}\,t^{17}\\ &- \frac{1}{1380784741023744000}\,t^{19} + \frac{1}{607545286050447360000}\,t^{21} + o(t^{-22})\end{aligned}$$

and

$$\begin{aligned}\mathbf{K}_{\mathrm{sech}}(0,t) = {}&-\frac{1}{4}\,t + \frac{1}{96}\,t^3 - \frac{1}{3840}\,t^5 + \frac{1}{258048}\,t^7 - \frac{1}{26542080}\,t^9 + \frac{1}{3892838400}\,t^{11}\\ &- \frac{1}{772905369600}\,t^{13} + \frac{1}{199766310912000}\,t^{15} - \frac{1}{65203723881676800}\,t^{17}\\ &+ \frac{1}{26234910079451136000}\,t^{19} - \frac{1}{12758451007059394560000}\,t^{21} + o(t^{-22})\end{aligned}$$

According to Theorem 8.7 we obtain the following table of the expansion coefficients for the integral kernel $\mathbf{K}_{\mathrm{sech}}$. Since all even coefficients are equal to 0, we do not include them in the table. To save the space, we represent all odd coefficients in the form of a fraction $\frac{x_n}{2^{n+1}n!}$.

n	1	3	5	7	9	11
b_n	$-\frac{1}{2^2 1!}$	$\frac{1}{2^4 3!}$	$-\frac{2}{2^6 5!}$	$\frac{5}{2^8 7!}$	$-\frac{14}{2^{10} 9!}$	$\frac{42}{2^{12} 11!}$
c_n	$-\frac{1}{2^2 1!}$	$\frac{3}{2^4 3!}$	$-\frac{10}{2^6 5!}$	$\frac{35}{2^8 7!}$	$-\frac{126}{2^{10} 9!}$	$\frac{462}{2^{12} 11!}$

n	13	15	17	19	21
b_n	$-\frac{132}{2^{14} 13!}$	$\frac{429}{2^{16} 15!}$	$-\frac{1430}{2^{18} 17!}$	$\frac{4862}{2^{20} 19!}$	$-\frac{16796}{2^{22} 21!}$
c_n	$-\frac{1716}{2^{14} 13!}$	$\frac{6435}{2^{16} 15!}$	$-\frac{24310}{2^{18} 17!}$	$\frac{92378}{2^{20} 19!}$	$-\frac{352716}{2^{22} 21!}$

For $f = \cosh x$ we have $h = f'(0) = 0$. The derivatives at the point $x = 0$ of the potential $q_1 \equiv 1$ are equal to $q_1^{(0)}(0) = 1$, $q_1^{(n)}(0) = 0$, $n \ge 1$, the non-zero values of the derivatives of the potential $q(x) = 1 - 2\,\mathrm{sech}^2(x)$ at the same point are given in the following table.

n	0	2	4	6	8	10	12
$q^{(n)}(0)$	-1	4	-32	544	-15872	707584	-44736512

n	14	16	18	20
$q^{(n)}(0)$	3807514624	-419730685952	58177770225664	-9902996106248192

The table below presents the values of the expression in parentheses appearing in (8.15), obtained from the derivatives of the potentials presented above with the coefficients

$S^n_{\ell;d;(n_1,\dots,n_\ell)}$ calculated by Proposition 8.3. All calculations were performed with the help of Matlab 7 and the results agree with the ones obtained from the explicit formulae for the integral kernels (see the table for b_n and c_n above). The value $n = 21$ is chosen only to keep the size of the table reasonable. Our Matlab program calculated exact values of the derivatives $\partial_t^n \mathbf{K}_{\cosh}(0,0)$ for $n \le 30$ and exact values of the derivatives $\partial_t^n \mathbf{K}_{\operatorname{sech}}(0,0)$ for $n \le 26$. To calculate the derivatives for larger values of n one have to use arbitrary precision arithmetic to overcome the limitation of the machine precision.

n	1	3	5	7	9	11
$2^{n+1}\partial_t^n \mathbf{K}_{\cosh}$	1	-3	10	-35	126	-462
$2^{n+1}\partial_t^n \mathbf{K}_{\operatorname{sech}}$	-1	1	-2	5	-14	42

n	13	15	17	19	21
$2^{n+1}\partial_t^n \mathbf{K}_{\cosh}$	1716	-6435	24310	-92378	352716
$2^{n+1}\partial_t^n \mathbf{K}_{\operatorname{sech}}$	-132	429	-1430	4862	-16796

9. *Approximate construction of integral kernels*

In this section we discuss two methods of approximate construction of integral kernels of transmutation operators. The first method is based on the expansion theorem and computation of the generalized Taylor coefficients by Theorem 8.7 and formula (8.15). The second method is based on the completeness result of hyperbolic formal powers and of generalized wave polynomials (Proposition 6.3 and [**18**, Theorem 22]) and on approximating the Goursat data for the integral kernel by generalized wave polynomials. However we present only the description of the main ideas of these approximate methods and do not perform detailed analysis. Such analysis, involving studying of approximative properties of the functions $\{u_n(x,x)\}$, convergency rate estimates and detailed comparison with existing numerical techniques, e.g., successive approximation and the series expansions from [**3**], goes beyond the scope of the present article. The authors hope to perform such analysis and present the results shortly.

The first method is applicable in the case when we know derivatives of the function $f \in C^n[-b,b]$ at the point $x = 0$ of all orders up to some order n, $n \ge 2$, i.e. when we know coefficients in the Taylor formula

$$f(x) = \sum_{k=0}^{n} f_k x^k + o(x^k).$$

Suppose additionally that the function f does not vanish on $[-b,b]$ and normalized as $f(0) = 1$, i.e. $f_0 = 1$. Then we can calculate Taylor coefficients up to the order n of the function $1/f$. For example, as it follows from [**10**, Theorem 11.7] or directly from the Faà di Bruno formula [**10**, Theorem 11.4],

$$\frac{1}{f(x)} = 1 + \sum_{k=1}^{n} \tilde{f}_k x^k + o(x^k),$$

where

$$\tilde{f}_k = \sum_{\substack{m_1+2m_2+\ldots+km_k=k\\ m_1\ge 0,\ldots,m_k\ge 0}} (-1)^{m_1+\ldots+m_k} \binom{m_1+\ldots+m_k}{m_1,\ldots,m_k} \prod_{j=1}^{k} f_j^{m_j}, \tag{9.1}$$

and $\binom{m_1+\ldots+m_k}{m_1,\ldots,m_k} = \frac{(m_1+\ldots+m_k)!}{m_1!\cdot\ldots\cdot m_k!}$ are multinomial coefficients.

Having coefficients f_k and $\tilde{f}_k$, $k \le n$, we easily calculate the Taylor coefficients up to the order $n-2$ of the potentials $q_f = f''/f$ and $q_{1/f} = f\cdot(1/f)''$. By (8.15) we obtain the values of the derivatives $\partial_t^m \mathbf{K}_f(0,0)$ and $\partial_t^m \mathbf{K}_{1/f}(0,0)$, $m \le n-1$, and by Theorem 8.7 we find the coefficients b_k and c_k, $k \le n-1$ in the representation (8.16). The obtained truncated series is an approximation of the integral kernel $\mathbf{K}_f$.

If on the opposite the Taylor coefficients of the potential q_f and the value of the parameter $h = f'(0)$ are known, the reconstruction of the Taylor coefficients of the solution f also presents no difficulty.

EXAMPLE 9.1. Consider the function $f = \cosh x$ as in Example 8.10. One can easily verify that the first Taylor coefficients for the function $1/f$ calculated by (9.1) coincide with the coefficients in the known expansion

$$\frac{1}{\cosh x} = \operatorname{sech} x = 1 - \frac{1}{2}x^2 + \frac{5}{24}x^4 - \frac{61}{720}x^6 + \frac{277}{8064}x^8 + O(x^{10}),$$

and the calculated Taylor coefficients for the potentials q_f and $q_{1/f}$ coincide with the values presented in Example 8.10.



FIGURE 1. *The integral kernel K_{sech} in the domain $\overline{\mathbf{R}}$ given by $b = 4$.*

We compare the exact integral kernel $\mathbf{K}_{\text{sech}}$ given by (8.21) with its approximations based on the generalized Taylor coefficients calculated in Example 8.10. To illustrate the dependence of the approximation error on the domain $\overline{\mathbf{R}}$ and on the number of coefficients used, we present the resulting approximation error for 3 different domains corresponding to values $b = 1$, $b = 2$

and $b = 4$, and for different number of generalized Taylor coefficients. The functions φ_k, defined by (2.23) and necessary for constructing the generalized wave polynomials, are calculated using two Matlab routines from the Spline Toolbox: on each step the integrand is approximated by a spline with 5000 knots using the command `spapi` and then it is integrated using `fnint`. The resulting kernel $\mathbf{K}_{\text{sech}}$ and its approximations were computed on the mesh of 100×100 equally spaced points in $\overline{\mathbf{R}}$. Obtained results together with the graphs of the integral kernel $\mathbf{K}_{\text{sech}}$ and the approximation error are given below.

$b = 1$		$b = 2$		$b = 4$	
N	Approx. error	N	Approx. error	N	Approx. error
1	0.12833	5	0.21204	13	1.8261
3	0.021458	7	0.043909	15	0.39987
5	0.0017866	9	0.0059553	17	0.070023
7	$8.9155 \cdot 10^{-5}$	11	0.00057228	19	0.010037
9	$2.9655 \cdot 10^{-6}$	13	$4.1076 \cdot 10^{-5}$	21	0.0011998
11	$7.0469 \cdot 10^{-8}$	15	$2.288 \cdot 10^{-6}$	23	0.00012146
13	$1.2562 \cdot 10^{-9}$	17	$1.0182 \cdot 10^{-7}$	25	$1.055 \cdot 10^{-5}$
15	$7.3683 \cdot 10^{-11}$	19	$3.7047 \cdot 10^{-9}$	27	$7.9493 \cdot 10^{-7}$
17	$7.3991 \cdot 10^{-11}$	21	$3.3373 \cdot 10^{-10}$	29	$5.2453 \cdot 10^{-8}$
19	$7.3989 \cdot 10^{-11}$	23	$3.3373 \cdot 10^{-10}$	31	$3.0562 \cdot 10^{-9}$



FIGURE 2. *The approximation error of the integral kernel* K_{sech} *in the domain* $\overline{\mathbf{R}}$ *given by* $b = 4$ *by the generalized wave polynomial of order* $N = 31$.

In order to estimate the error of approximation the exact formula (8.21) for the kernel $\mathbf{K}_{\text{sech}}$ was used, where the corresponding integrals were computed numerically. As can be seen from (8.21) the integrands for t close to x involve subtraction of close values and division by values close to zero. Due to this fact the kernel $\mathbf{K}_{\text{sech}}$ computed in this way by (8.21) is not exact. This explains why in the above tables after a certain value of N the reported approximation error does not decrease, it is limited by the computational error of $\mathbf{K}_{\text{sech}}$.

To the difference of $\mathbf{K}_{\text{sech}}$ the integral kernel $\mathbf{K}_{\text{cosh}}$ does not involve integration and may be computed more precisely. The approximations of the integral kernel $\mathbf{K}_{\text{cosh}}$ were computed for $b = 2$ and the resulting absolute errors obtained for different numbers of expansion coefficients are presented in the table below.

$b = 2$					
N	Approx. error	N	Approx. error	N	Approx. error
1	1.7878	11	0.00074903	21	$1.316 \cdot 10^{-10}$
3	1.088	13	$5.2024 \cdot 10^{-5}$	23	$3.3386 \cdot 10^{-12}$
5	0.33724	15	$2.8243 \cdot 10^{-6}$	25	$7.5051 \cdot 10^{-14}$
7	0.063779	17	$1.2312 \cdot 10^{-7}$	27	$6.9944 \cdot 10^{-15}$
9	0.0081416	19	$4.4042 \cdot 10^{-9}$	29	$6.6613 \cdot 10^{-15}$

The first method of approximate construction of the integral kernels requires the knowledge of higher order derivatives of the potentials q_f and $q_{1/f}$ in the origin. In the case when such derivatives are unknown or hard to obtain analytically, instead of performing numerical differentiation we can approximate the integral kernel by approximating its Goursat data given by (3.6) with truncated series of the form (4.3) and (4.4).

Indeed, suppose that the numbers $c_0, \dots, c_N$ and $b_1, \dots, b_N$ are such that

$$\left|\frac{1}{2}\big(\mathbf{K}_f(x,x) + \mathbf{K}_f(x,-x)\big) - c_0 u_0(x,x) - \sum_{n=1}^{N} c_n u_{2n-1}(x,x)\right| < \varepsilon_1$$

and

$$\left|\frac{1}{2}\big(\mathbf{K}_f(x,x) - \mathbf{K}_f(x,-x)\big) - \sum_{n=1}^{N} b_n u_{2n}(x,x)\right| < \varepsilon_2$$

for every $x \in [-b, b]$. Consider the functions

$$K(x,t) = c_0 u_0(x,t) + \sum_{n=1}^{N} c_n u_{2n-1}(x,t) + \sum_{n=1}^{N} b_n u_{2n}(x,t) \tag{9.2}$$

and $\widetilde{\mathbf{K}}_f = T_f^{-1}\mathbf{K}_f$ and $\widetilde{K} = T_f^{-1}K$. Then by the definition of the Goursat-to-Goursat transmutation operator

$$\begin{pmatrix} \widetilde{\mathbf{K}}_f(x,x) \\ \widetilde{\mathbf{K}}_f(x,-x) \end{pmatrix} = T_G^{-1} \begin{pmatrix} \mathbf{K}_f(x,x) \\ \mathbf{K}_f(x,-x) \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} \widetilde{K}(x,x) \\ \widetilde{K}(x,-x) \end{pmatrix} = T_G^{-1} \begin{pmatrix} K(x,x) \\ K(x,-x) \end{pmatrix},$$

hence due to the boundedness of the operator T_G^{-1}

$$\max\Big(\max_{x\in[-b,b]}\big|\widetilde{\mathbf{K}}_f(x,x) - \widetilde{K}(x,x)\big|, \max_{x\in[-b,b]}\big|\widetilde{\mathbf{K}}_f(x,-x) - \widetilde{K}(x,-x)\big|\Big)$$

$$\begin{aligned}&\le \|T_G^{-1}\| \max\Big(\max_{x\in[-b,b]} \big|\mathbf{K}_f(x,x) - K(x,x)\big|, \max_{x\in[-b,b]} \big|\mathbf{K}_f(x,-x) - K(x,-x)\big|\Big)\\ &\quad\le \|T_G^{-1}\| \max_{x\in[-b,b]} \bigg(\left|\frac{1}{2}\big(\mathbf{K}_f(x,x)+\mathbf{K}_f(x,-x)\big) - \frac{1}{2}\big(K(x,x)+K(x,-x)\big)\right| \\ &\qquad + \left|\frac{1}{2}\big(\mathbf{K}_f(x,x)-\mathbf{K}_f(x,-x)\big) - \frac{1}{2}\big(K(x,x)-K(x,-x)\big)\right|\bigg) < \|T_G^{-1}\|(\varepsilon_1+\varepsilon_2),\end{aligned}$$

where we have used equalities $\frac{1}{2}\big(K(x,x)+K(x,-x)\big) = c_0u_0(x,x) + \sum_{n=1}^N c_n u_{2n-1}(x,x)$ and $\frac{1}{2}\big(K(x,x)-K(x,-x)\big) = \sum_{n=1}^N b_n u_{2n}(x,x)$. We obtain from the proof of [**18**, Theorem 3] that for every $(x,t)\in\overline{\mathbf{R}}$

$$\big|\widetilde{\mathbf{K}}_f(x,t) - \widetilde{K}(x,t)\big| \le 3\|T_G^{-1}\|(\varepsilon_1+\varepsilon_2),$$

hence for every $(x,t)\in\overline{\mathbf{R}}$

$$\big|\mathbf{K}_f(x,t) - K(x,t)\big| \le 3\|T_f\|\cdot\|T_G^{-1}\|(\varepsilon_1+\varepsilon_2).$$

That is, if we approximate the function $g_1(x) := \frac{1}{2}\big(\mathbf{K}_f(x,x)+\mathbf{K}_f(x,-x)\big) = \frac{h}{2} + \frac{1}{4}\int_0^x q_f(s)\,ds$ by the functions $\{u_0(x,x)\}\cup\{u_{2n-1}(x,x)\}_{n\ge1}$ and the function $g_2(x) := \frac{1}{2}\big(\mathbf{K}_f(x,x) - \mathbf{K}_f(x,-x)\big) = \frac{1}{4}\int_0^x q_f(s)\,ds$ by the functions $\{u_{2n}(x,x)\}_{n\ge1}$, we obtain an approximation having the form (9.2) of the integral kernel. It is worth to mention that we do not need to know the Darboux associated potential $q_{1/f}$ for this method and we do not impose additional smoothness conditions on the potential q_f. The approximation of the functions g_1 and g_2 by the corresponding combinations of the functions $u_n(x,x)$ can be done in several ways. We may use the least squares method to obtain a good, however far from the best, approximation. We may reformulate the approximation problem as a linear programming problem and solve it. Under the additional assumption that the corresponding systems of the functions $u_n(x,x)$ satisfy the Haar condition, we may use the Remez algorithm. We refer the reader to [**18**, Section 6] and to [**35**, §7], [**39**, Chapter 6] and [**40**, Chapter 1] for further details.

Example 9.2. We approximated the integral kernels $\mathbf{K}_{\text{sech}}$ and $\mathbf{K}_{\text{cosh}}$ from Example 9.1 by the second described method. All the parameters are taken as in the previous example. The Goursat data were approximated by corresponding functions $u_n(x,x)$ with the help of the Remez simple exchange algorithm, see e.g., [**35**, §7].

$b=2$, $\mathbf{K}_{\text{sech}}$ kernel	
N	Approx. error
5	0.0045993
9	$9.3687\cdot10^{-6}$
13	$4.1549\cdot10^{-9}$
17	$3.3354\cdot10^{-10}$

$b=4$, $\mathbf{K}_{\text{sech}}$ kernel	
N	Approx. error
13	$7.4042\cdot10^{-5}$
17	$2.342\cdot10^{-7}$
21	$2.789\cdot10^{-10}$
25	$4.9467\cdot10^{-11}$

$b=2$, $\mathbf{K}_{\text{cosh}}$ kernel	
N	Approx. error
5	0.0052907
9	$1.2563\cdot10^{-5}$
13	$6.6227\cdot10^{-9}$
17	$1.1813\cdot10^{-12}$
19	$1.0325\cdot10^{-14}$

References

1. H. Begehr and R. Gilbert, *Transformations, transmutations and kernel functions, vol. 1–2* (Longman Scientific & Technical, Harlow, 1992).

2. L. Bers, *Theory of pseudo-analytic functions* (New York University, 1952).

3. A. Boumenir, ‘The approximation of the transmutation kernel’, *J. Math. Phys.* 47 (2006) 013505.



FIGURE 3. *The approximation error of the integral kernel* $K_{\cosh}$ *in the domain* $\overline{\mathbf{R}}$ *given by* $b = 2$ *by the generalized wave polynomial of order* $N = 19$. *The generalized wave polynomial is obtained with the use of the Remez simple exchange algorithm.*

4. H. CAMPOS and V. V. KRAVCHENKO, 'Fundamentals of bicomplex pseudoanalytic function theory: Cauchy integral formulas, negative formal powers and Schrödinger equations with complex coefficients', *Complex Analysis and Operator Theory* (2012) DOI: 10.1007/s11785-012-0256-4.

5. H. M. CAMPOS, V. V. KRAVCHENKO and L. M. MENDEZ, 'Complete families of solutions for the Dirac equation: an application of bicomplex pseudoanalytic function theory and transmutation operators', *Advances in Applied Clifford Algebras* (2012) DOI: 10.1007/s00006-012-0349-1.

6. H. CAMPOS, V. V. KRAVCHENKO and S. M. TORBA, 'Transmutations, L-bases and complete families of solutions of the stationary Schrödinger equation in the plane', *J. Math. Anal. Appl.* 389 (2012) 1222–1238.

7. R. W. CARROLL, *Transmutation theory and applications* (Mathematics Studies, Vol. 117, North-Holland, 1985).

8. A. CASTAÑEDA and V. V. KRAVCHENKO, 'New applications of pseudoanalytic function theory to the Dirac equation', *J. Phys. A: Math. Gen.* 38 (2005) 9207–9219.

9. R. CASTILLO, V. V. KRAVCHENKO, H. OVIEDO and V. S. RABINOVICH, 'Dispersion equation and eigenvalues for quantum wells using spectral parameter power series', *J. Math. Phys.* 52 (2011) 043522 (10 pp.).

10. CH. A. CHARALAMBIDES, *Enumerative combinatorics* (Chapman & Hall/CRC, Boca Raton, 2002).

11. L. COMTET, *Advanced combinatorics* (D. Reidel, Dordrecht, Holland, 1974).

12. J. DELSARTE, 'Sur certaines transformations fonctionnelles relatives aux équations linéaires aux dérivées partielles du second ordre', *C. R. Acad. Sc.* 206 (1938) 178–182.

13. J. DELSARTE and J. L. LIONS, 'Transmutations d'opérateurs différentiels dans le domaine complexe', *Comment. Math. Helv.* 32 (1956) 113–128.

14. L. ERBE, R. MERT and A. PETERSON 'Spectral parameter power series for Sturm–Liouville equations on time scales', *Applied Mathematics and Computation* 218 (2012) 7671–7678.

15. M. K. FAGE and N. I. NAGNIBIDA, *The problem of equivalence of ordinary linear differential operators* (Novosibirsk: Nauka, 1987) (in Russian).

16. T. KATO, *Perturbation theory for linear operators* (Springer-Verlag, 1966).

17. K. V. KHMELNYTSKAYA, V. V. KRAVCHENKO and H. C. ROSU, 'Eigenvalue problems, spectral parameter power series, and modern applications', Submitted, available at arXiv:1112.1633.

18. K. V. KHMELNYTSKAYA, V. V. KRAVCHENKO, S. M. TORBA and S. TREMBLAY, 'Wave polynomials, transmutations and Cauchy's problem for the Klein-Gordon equation', Submitted, available at arXiv:1208.5984.

19. K. V. KHMELNYTSKAYA and H. C. ROSU, 'A new series representation for Hill's discriminant', *Annals of Physics* 325 (2010) 2512-2521.

20. V. G. KRAVCHENKO, V. V. KRAVCHENKO and S. TREMBLAY, 'Zakharov-Shabat system and hyperbolic pseudoanalytic function theory', *Math. Meth. Appl. Sci.* 33 (2010) 448–453.

21. V. V. KRAVCHENKO, 'On a relation of pseudoanalytic function theory to the two-dimensional stationary

Schrödinger equation and Taylor series in formal powers for its solutions', *J. Phys. A: Math. Gen.* 38 (2005) 3947–3964.

22. V. V. Kravchenko, 'A representation for solutions of the Sturm-Liouville equation', *Complex Variables and Elliptic Equations* 53 (2008) 775–789.

23. V. V. Kravchenko, *Applied pseudoanalytic function theory* (Basel: Birkhäuser, Series: Frontiers in Mathematics, 2009).

24. V. V. Kravchenko and R. M. Porter, 'Spectral parameter power series for Sturm-Liouville problems', *Math. Meth. Appl. Sci.* 33 (2010) 459–468.

25. V. V. Kravchenko, D. Rochon and S. Tremblay, 'On the Klein-Gordon equation and hyperbolic pseudoanalytic function theory', *J. Phys. A: Math. Gen.* 41 (2008) 065205, (18pp.).

26. V. V. Kravchenko and S. Torba, 'Transmutations for Darboux transformed operators with applications', *J. Phys. A: Math. Gen.* 45 (2012) 075201 (21 pp.).

27. V. V. Kravchenko and S. M. Torba 'Transmutations and spectral parameter power series in eigenvalue problems', To appear in *Operator Theory: Advances and Applications*, available at arXiv:1203.4225.

28. V. V. Kravchenko and S. M. Torba, 'Spectral problems in inhomogeneous media, spectral parameter power series and transmutation operators', *Proceedings of the 14-th International Conference on Mathematical Methods in Electromagnetic Theory, August 2012 – Kharkiv, Ukraine*, 18–22. Available at arXiv:1207.2713.

29. V. V. Kravchenko and S. Tremblay, 'Explicit solutions of generalized Cauchy-Riemann systems using the transplant operator', *J. Math. Anal. Appl.* 370 (2010) 242–257.

30. G. L. Lamb, *Elements of soliton theory* (John Wiley & Sons, New York, 1980).

31. M. A. Lavrentyev and B. V. Shabat, *Hydrodynamics problems and their mathematical models* (Nauka, Moscow, 1977) (in Russian).

32. B. M. Levitan, *Inverse Sturm-Liouville problems* (VSP, Zeist, 1987).

33. J. L. Lions, 'Solutions élémentaires de certains opérateurs différentiels à coefficients variables', *Journ. de Math.* 36 (1957) 57–64.

34. V. A. Marchenko, *Sturm-Liouville operators and applications* (Basel: Birkhäuser, 1986).

35. G. Meinardus, *Approximation of Functions: Theory and Numerical Methods* (New York: Springer, 1967).

36. A. F. Motter and M. A. F. Rosa, 'Hyperbolic calculus', *Adv. Appl. Clifford Algebras* 8 (1998) 109–128.

37. N. A. Pakhareva and I. N. Aleksandrovich, 'Representation of $p_1(x)p_2(y)$-wave functions by a linear combination of wave functions and their derivatives' (Russian), *Dokl. Akad. Nauk Ukrain. SSR Ser. A* 1983 no. 1, 23–27.

38. N. A. Pakhareva and I. N. Aleksandrovich, 'Integral representation of p-wave functions with characteristic $p = e^{\alpha x}y^k$' (Russian), *Vychisl. Prikl. Mat.(Kiev)* 49 (1983) 35–42.

39. J. R. Rice, *The Approximation of Functions, Vol. 1. Linear theory* (Reading, Massachusetts: Addison-Wesley, 1964).

40. T. J. Rivlin, *An introduction to the approximation of functions* (Blaisdell: Waltham, Mass., 1969).

41. D. Rochon and S. Tremblay, 'Bicomplex quantum mechanics: I. The generalized Schrödinger equation', *Advances in Applied Clifford Algebras* 14 (2004) 231–248.

42. S. M. Sitnik, *Transmutations and applications: a survey.* arXiv:1012.3741v1, originally published in the book: *Advances in Modern Analysis and Mathematical Modeling* (Editors: Yu.F.Korobeinik, A.G.Kusraev, Vladikavkaz: Vladikavkaz Scientific Center of the Russian Academy of Sciences and Republic of North Ossetia–Alania, 2008, 226–293).

43. G. Sobczyk, 'The hyperbolic number plane', *Coll. Maths. Jour.* 26 (1995) 268–280.

44. K. Trimeche, *Transmutation operators and mean-periodic functions associated with differential operators* (London: Harwood Academic Publishers, 1988).

45. I. N. Vekua, *Generalized analytic functions* (Moscow: Nauka, 1959 (in Russian); English translation Oxford: Pergamon Press 1962).

46. V. S. Vladimirov, *Equations of mathematical physics* (Moscow: Nauka, 1984 (in Russian); Engl. transl.: of the first edition: N.Y.: Marcel Dekker, 1971).

47. G. C. Wen, *Linear and Quasilinear Complex Equations of Hyperbolic and Mixed Type* (Taylor & Francis London, 2003).

Vladislav V. Kravchenko
Department of Mathematics,
CINVESTAV del IPN,
Unidad Queretaro,
Libramiento Norponiente No. 2000,
Fracc. Real de Juriquilla, Queretaro,
Qro. C.P. 76230 MEXICO

vkravchenko@math.cinvestav.edu.mx

Sergii M. Torba
Department of Mathematics,
CINVESTAV del IPN,
Unidad Queretaro,
Libramiento Norponiente No. 2000,
Fracc. Real de Juriquilla, Queretaro,
Qro. C.P. 76230 MEXICO

storba@math.cinvestav.edu.mx